

# Discovery of an Excess of H $\alpha$ Emitters around 4C 23.56 at $z = 2.48$

Ichu TANAKA\*, <sup>1</sup> Carlos DE BREUCK, <sup>2</sup> Jaron D. KURK, <sup>3</sup> Yoshiaki TANIGUCHI, <sup>4</sup>  
Tadayuki KODAMA, <sup>1,5</sup> Yuichi MATSUDA, <sup>6</sup> Chris PACKHAM, <sup>7</sup> Andrew ZIRM, <sup>8</sup> Masaru KAJISAWA, <sup>5,9</sup>  
Takashi ICHIKAWA, <sup>9</sup> Nick SEYMOUR, <sup>10</sup> Daniel STERN, <sup>11</sup> Alan STOCKTON, <sup>12</sup> Bram P. VENEMANS, <sup>2</sup>  
and

Joël VERNET <sup>2</sup>

<sup>1</sup>*Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'ohoku Place, Hilo, HI 96720, USA*

<sup>2</sup>*European Southern Observatory, Karl-Schwarzschild Strasse 2, 85748 Garching bei München, Germany*

<sup>3</sup>*Max-Planck-Institut für Extraterrestrische Physik, Postfach 1312, 85741 Garching, Germany*

<sup>4</sup>*Research Center for Space and Cosmic Evolution, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan*

<sup>5</sup>*National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*

<sup>6</sup>*Department of Physics, University of Durham, South Road, Durham DH1 3LE, UK*

<sup>7</sup>*Department of Astronomy, University of Florida, 211 Bryant Science Center, P.O. Box 112055, Gainesville, FL 32611-2055, USA*

<sup>8</sup>*Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen, Denmark*

<sup>9</sup>*Astronomical Institute, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan*

<sup>10</sup>*Mullard Space Science Laboratory, UCL, Holmbury St Mary, Dorking, Surrey, RH5 6NT, UK*

<sup>11</sup>*Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109, USA*

<sup>12</sup>*Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA*

(Received ; accepted )

## Abstract

We report the discovery of a significant excess of candidate H $\alpha$  emitters (HAEs) in the field of the radio galaxy 4C 23.56 at  $z = 2.483$ . Using the MOIRCS near-infrared imager on the Subaru Telescope we found 11 candidate emission-line galaxies to a flux limit of  $\sim 7.5 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ , which is about 5 times excess from the expected field counts with  $\sim 3\text{-}\sigma$  significance. Three of these are spectroscopically confirmed as redshifted H $\alpha$  at  $z = 2.49$ . The distribution of candidate emitters on the sky is tightly confined to a 1.2-Mpc-radius area at  $z = 2.49$ , locating 4C 23.56 at the western edge of the distribution. Analysis of the deep *Spitzer* MIPS 24  $\mu\text{m}$  imaging shows that there is also an excess of faint MIPS sources. All but two of the 11 HAEs are also found in the MIPS data. The inferred star-formation rate (SFR) of the HAEs based on the extinction-corrected H $\alpha$  luminosity (median SFR  $\gtrsim 100 \text{ M}_{\odot} \text{ yr}^{-1}$ ) is similar to those of HAEs in random fields at  $z \sim 2$ . On the other hand, the MIPS-based SFR for the HAEs is on average 3.6 times larger, suggesting the existence of the star-formation significantly obscured by dust. The comparison of the H $\alpha$ -based star-formation activities of the HAEs in the 4C 23.56 field to those in another proto-cluster around PKS 1138-262 at  $z = 2.16$  reveals that the latter tend to have fainter H $\alpha$  emission despite similar  $K$ -band magnitudes. This suggests that star-formation may be suppressed in the PKS 1138-262 protocluster relative to the 4C 23.56 protocluster. This difference among the HAEs in the two proto-clusters at  $z > 2$  may imply that some massive cluster galaxies are just forming at these epochs with some variation among clusters.

**Key words:** Galaxies: protocluster — Galaxies: individual (4C 23.56) — Galaxies: high redshifts — Galaxies: H $\alpha$  emitters

## 1. Introduction

Clusters of galaxies are the most massive bound systems in the universe. The present-day cluster population is dominated by early-type galaxies which make up a tight “red sequence” in color-magnitude diagram (Dressler 1980). The analyses of the zero-point, slope, and scatter of the color-magnitude relation have been playing key roles in understanding when and how galaxies in clusters are

formed and evolved (e.g., Bower Lucey, & Ellis 1992).

Studies of low to intermediate redshift clusters indicate that the majority of stars in cluster galaxies are formed at  $z > 2$  (e.g., Ellis et al. 1997; Bower et al. 1998; Kodama et al. 1998; Stanford et al. 1998; Holden et al. 2004; Tran et al. 2007). On the other hand, detailed morphological analyses of the cluster red galaxies suggest that a significant fraction of the present-day cluster early-type galaxies must have been fully assembled after  $z \sim 1$  (e.g., Dressler et al. 1997; van Dokkum et al. 2000; van Dokkum & Franx 2001; Ellis et al. 2006; Mei et al. 2009). A significant rise in

\* ichi@subaru.aoj.org

star-formation activity in the cluster cores is also seen as one goes to higher ( $z > 0.4$ ) redshift, the “Butcher-Oemler effect” (Butcher & Oemler 1978; Butcher & Oemler 1984).

Studies of the Butcher-Oemler effect, as well as the morphological transformation in cluster cores, have been among the major topics in extragalactic astronomy for the past two decades (e.g., Dressler et al. 1994; Couch et al. 1994; Dressler et al. 1997; Balogh et al. 1997; Oemler et al. 1997; Poggianti et al. 1999; van Dokkum et al. 2000; Ellingson et al. 2001; Kodama & Bower 2001; Mei et al. 2009; Pannella et al. 2009). As the observational data have been accumulated for ever more distant galaxy clusters, the “break” of the red sequence at fainter magnitudes and its evolution has attracted great attention (e.g., Kajisawa et al. 2000; Kodama et al. 2004; De Lucia et al. 2004; Tanaka et al. 2005; De Lucia et al. 2007; Gilbank et al. 2008). These phenomena have been discussed in the framework of the “down-sizing” of the star-formation activity, whereby the stellar mass of the galaxies which host star formation at a fixed rate is, on average, decreasing with cosmic time. The evolution of the faint end of the red sequence is also connected to the continuous infall of star-forming galaxies into the cluster cores, their subsequent cessation of star-formation, and change in their morphologies (Poggianti et al. 2004; Bundy et al. 2006).

From the local universe to intermediate redshifts ( $z < 1$ ), a number of studies suggest that star formation is strongly suppressed in the cluster cores compared with the low-density field environment (e.g., Hashimoto et al. 1998; Gómez et al. 2003; Nakata et al. 2005; Poggianti et al. 2006; Patel et al. 2009). For passively-evolving early-type galaxies, there is also weak, but growing, evidence that the luminosity-weighted ages of galaxies tend to be older in higher density environments. This suggests that perhaps the time scale of star formation and/or secondary episodes of the star formation vary with environment (Thomas et al. 2005; Clemens et al. 2006; Rakos et al. 2008; Clemens et al. 2009; Blanton & Moustakas 2009; Pannella et al. 2009; Thomas et al. 2010). Recently, however, this trend is observed to flatten or even reverse, with the density dependence of star-formation disappearing or with enhanced star-formation in the cluster core at  $z > 1$  (Elbaz et al. 2007; Ideue et al. 2009; Hayashi et al. 2010; Tran et al. 2010).

If the majority of the massive cluster galaxies ( $M_* > 10^{10} M_\odot$ ) begin forming their stars at  $z > 2$ , they will have high star formation rates at higher redshifts. Hayashi et al. (2010) have surveyed star-forming galaxies in the X-ray cluster XMMXCS J2215.9-1738 at  $z = 1.46$  using a narrowband filter that samples the redshifted [O II] doublet located at the cluster redshift. They showed that the [O II] emitting star-forming galaxies are prevalent everywhere in the cluster including its core region (see also, Hilton et al. 2010). Although star formation is still confined to the cluster outskirts in other  $z \sim 1.4$  cluster (Lidman et al. 2008; Bauer et al. 2010; Strazzullo et al. 2010), the widespread star formation in XMMXCS J2215.9-1738 is in strong contrast to similar surveys of lower redshift clusters at  $z = 0.81$  and  $z = 0.41$  (Kodama

et al. 2004; Koyama et al. 2010). The color-magnitude relation for XMMXCS J2215.9-1738 also shows a larger scatter than those of other intermediate redshift clusters (Hilton et al. 2009; Papovich et al. 2010). Although we need a larger sample of clusters at intermediate and high redshifts to discern a general trend in the star formation – galaxy density relation (Elbaz et al. 2007; Poggianti et al. 2008; Sobral et al. 2010), the existence of widespread star formation activity in this  $z = 1.5$  cluster strongly suggests that we are gradually approaching the major epoch of galaxy formation in cluster environments.

At higher redshifts ( $z > 2$ ), the number of the known (proto-) clusters is still quite small, and the majority of the known proto-clusters were discovered in narrow-band Ly $\alpha$ -emitter surveys. With some exceptions (e.g., Steidel et al. 1998; Steidel et al. 2000; Shimasaku et al. 2003; Steidel et al. 2005; Gobat et al. 2010), much of the search for proto-clusters has been done around powerful radio galaxies and QSOs with known redshifts (e.g., Keel et al. 1999; Pentericci et al. 2000; Kurk et al. 2000; Yamada et al. 2001; Venemans et al. 2002; Kurk et al. 2004a; Venemans et al. 2005; Venemans et al. 2007; Matsuda et al. 2009). The success of these searches around radio galaxies is related to the fact that the host galaxies of the powerful radio sources tend to be the most massive galaxies at any epoch of the universe (De Breuck et al. 2002; Seymour et al. 2007). Powerful radio sources are powered by central super-massive black holes, and the large inferred masses of the black holes in turn imply massive host stellar systems (Magorrian et al. 1998). The largest galaxies in the universe tend to lie in the centers of largest-scale haloes which must have originated from the primordial density peaks where galaxy formation processes are accelerated in a biased way (Rees 1985).

Recently Tanaka et al. (2010) analyzed the broadband SED of galaxies in the PKS 1138 proto-cluster and found that proto-cluster galaxies show suppressed star-formation on average compared to field galaxies (see also, Zirm et al. 2008; Doherty et al. 2010). This implies that even at  $z \gtrsim 2$  the activity in the cluster core region seems to lag behind the star-formation activity we see in galaxies in field surveys. Steidel et al. (2000) also reported older stellar ages for their  $z = 2.3$  proto-cluster than for field galaxies in the same survey, implying a similar trend. The detection of a possible (proto-) red-sequence in four known proto-cluster fields at  $z \gtrsim 2$  via *JHK* photometry (Kodama et al. 2007) also indicates that, though the star-formation is also likely to be prevalent there, some *K*-bright galaxies may already be converging to the red-sequence.

At higher redshifts, there are some hints that more intense star-formation is occurring even in the proto-cluster environment. In the SSA22 field, which is known to contain a large overdensity of galaxies at  $z = 3.1$  (Steidel et al. 1998; Steidel et al. 2000), Matsuda et al. (2004) detected a significant number of the extended Ly $\alpha$  blobs which are thought to be progenitors of massive galaxies (e.g., Dijkstra & Loeb 2009). Two of the largest Ly $\alpha$  blobs are identified with red, massive ( $M_* \sim 10^{11} M_\odot$ ) galax-

ies, comparable to dominant cluster galaxies at  $z < 1.5$  (Uchimoto et al. 2008; Stott et al. 2010). Tamura et al. (2009) have detected an excess of AzTEC 1.1 mm-sources in the same SSA22 field. The positional correlation of the AzTEC sources with the Ly $\alpha$  emitter distribution at  $z = 3.1$  suggests that dust-enshrouded vigorous star-formation might be going on in the field. There is also a report of a significant excess of submm-bright objects in the TNJ 1338-1942 field at  $z = 4.1$ , which is known to have a proto-cluster identified by Ly $\alpha$  emitters (De Breuck et al. 2004; Venemans et al. 2007).

As was demonstrated by Kodama & Bower (2001), star formation activity in cluster cores should have been as vigorous as in the field environment at some point in the past (and may exceed it at even higher redshifts). To observe this crossover point we need larger samples of proto-clusters at  $z = 2-4$ .

The field around 4C 23.56 is one of the most promising proto-cluster candidates at  $z > 2$  (Knopp & Chambers 1997, hereafter KC97). The authors detected an excess of candidate Ly $\alpha$  emitters as well as an excess of very red objects in  $I - K'$  color in their survey field ( $1.25 \text{ arcmin}^2$ ). More recently, Kajisawa et al. (2006a) took deep  $J$  and  $K_s$  images with CISCO on Subaru and found a significant excess of Distant Red Galaxies (DRGs, Franx et al. 2003) in a  $1.6 \times 1.6 \text{ arcmin}^2$  field. This further supports the idea that there are a number of evolved galaxies in this proto-cluster at  $z \sim 2.49$ . However, both of these surveys have very limited field coverage and they may not be probing the whole picture of the proto-cluster. So far no other follow-up observations, including any redshift confirmation, have been reported in the literature.

Here we present the result of our new H $\alpha$  emitter (HAE) search around 4C 23.56 with the wide-field near-infrared imager, MOIRCS, mounted on the 8.2m Subaru Telescope. In this study, we show that there is a clear and strong excess of the HAEs near the radio galaxy. We present their star formation properties and discuss the characteristics of this proto-cluster by comparing with other fields at similar redshifts. Archival *Spitzer* MIPS data are also used to probe dusty star forming activity in the proto-cluster. This is the first narrow-band H $\alpha$  emitter survey of a proto-cluster field at  $z > 2$  after Kurk et al. (2004a). The field size we cover is nearly 19 times larger than that of KC97, and is more than twice as large as that of Kurk et al. (2004a). We note that the survey redshift of  $z \sim 2.5$  is close to the red end of the H $\alpha$ -observable window ( $z < 2.7$ ) from the ground, due to the significant rise of the thermal background beyond the  $K$ -band window.

Throughout the paper, the standard  $\Lambda$ CDM cosmology with  $\Omega_m = 0.3, \Omega_\Lambda = 0.7, h = 0.7$  is assumed ( $1' = 1.18$  comoving Mpc at  $z = 2.48$ ). All the magnitudes presented in this paper are in the AB system unless otherwise noted.

## 2. Observations and Data Reduction

### 2.1. Observations

The primary data for the current work were collected using the Multi-Object Infrared Camera and

Spectrograph (MOIRCS, Ichikawa et al. 2006; Suzuki et al. 2008) mounted on the Cassegrain focus of the Subaru 8.2-m Telescope at Mauna Kea. The observations were executed during the second half-night of UT 1st June, 2007, during the author's (IT) Subaru Staff Time. The weather was photometric but with variable seeing in the range  $0''.47 - 0''.7$ .

MOIRCS has two independent sets of optics (including both filters and detectors), which divide the full  $4' \times 7'$  field of view (FOV) into the two subfields (channels 1 and 2), each covering  $4' \times 3'.5$ . We set the MOIRCS position angle to  $-90^\circ$  so that the long side ( $= 7'$  side) of the FOV aligns to the east-west direction. In this setting, the channel 1 FOV is on the eastern (left) side (see Fig.1b).

We used both the standard  $K_s$ -band filter and a narrow-band (NB) filter called “CO” or “NB2288”. The NB filter has central wavelength  $\lambda_c = 2.2881 \mu\text{m}$  with a bandwidth of  $0.023 \mu\text{m}$ . Note that these are the post-installation measured values (M. Akiyama, private communication). Individual NB exposures were 300 s. In total, we obtained 16 exposures, yielding a total integration time of 4800 s. Shallow data were also acquired in the  $K_s$  band to enable continuum subtraction of the NB data. The  $K_s$  band exposures were taken in 55 s and 3-coadds. We obtained 9 frames in total (1485 s). We found that stars on three images were elongated due to poor guiding at high elevation ( $> 84$  degrees). We rejected those frames during the final combine, resulting in a total integration of 990 s. We applied a standard circular-dither pattern with a diameter of  $20''$ . After the target observation, four position dithered data of the FS151 standard star were acquired in both the  $K_s$  and NB2288 bands.

We note that 4C 23.56 is close ( $\sim 1^\circ$ ) to a bright ( $K \sim 0$  mag) Mira-type star. Such bright stars are known to occasionally cause significant stray light on the image<sup>1</sup>. Fortunately, the effect was small during the night. However, stray light was seen in some of the data. It affected only a tiny portion ( $< 1\%$ ) of the total area, and we took care to mark the positions of any stray light before identifying the emitters. Unfortunately, stray light caused a hole-like pattern near the radio galaxy in the  $K_s$  data. Due to this, the total magnitude for 4C 23.56 may be systematically wrong. For other targets, we confirmed that stray light does not affect the photometry.

#### 2.1.1. Wavelength Shift of NB filters

In general, the central wavelength of NB filters will shift with the angle of incidence of the light to the filter surface. In our case, therefore, the observable wavelength window will change with the position of targets on the detector.

In Figure 1 we show the simulated shift of the filter transmission curve with source position on the detector (the case for channel 2 is shown here: R. Suzuki, private comm.). We find that the redshift range of interest,  $2.47 < z < 2.49$ , is always covered by  $> 80\%$  of the detector area. As a reference, the expected wavelength of the H $\alpha$  emission from 4C 23.56 is plotted as the arrow in Fig.1. Note that the redshift of 4C 23.56 is  $2.483 \pm 0.003$

<sup>1</sup> See the *Imaging Information* page on MOIRCS website.

by Röttgering et al. (1997) from C IV, He II, and C III] lines. The bottom of Fig.1 shows the iso-redshift lines of the observable H $\alpha$  emission for the center of the filter transmission. The iso-redshift line for 2.483 goes across the center of the detector. In the upper panel of Fig.1 we show that H $\alpha$  emission at  $\pm 500$  km s $^{-1}$  offset from 4C 23.56 can be observed everywhere but the very edge of the detector. The plot shows that the emitters blueshifted by  $\lesssim 1000$  km s $^{-1}$  relative to the radio galaxy will leave the filter at the center of the FOV, while for objects redshifted by  $\gtrsim 1000$  km s $^{-1}$  escape our detection in the outer 1/4 of the detector. Though MOIRCS uses two independent filters for the two channels, the difference of the center wavelength of these two NBs is only 3 Å ( $\sim 40$  km s $^{-1}$  difference at  $z = 2.49$ ). It should not cause any systematics between the channels.

We note that the velocity window of  $\pm 500$  km s $^{-1}$  is just adequate for cluster member selection. Simulations of X-ray clusters shows that the velocity dispersion of rich clusters generally gets smaller by about a factor of 2 as one goes from  $z = 0$  to 3 (e.g., Eke et al. 1998). Also, the previous H $\alpha$  emitter survey for a dense protocluster at  $z = 2.16$  by Kurk et al. (2004b) found a velocity dispersion of only 360 km s $^{-1}$  (narrower than the width of the NB filter for their pre-selection), small for the inferred richness of the protocluster. Thus we would expect to detect a significant fraction of the protocluster H $\alpha$  emitters within the MOIRCS narrowband window if its dynamical center is at the same redshift and located on 4C 23.56.

## 2.2. MOIRCS Data Reduction by MCSRED

Data reduction of the MOIRCS science frames was performed using the MCSRED package<sup>2</sup>, the IRAF<sup>3</sup> script pipeline for MOIRCS imaging data developed by the author (IT). Here we briefly describe the procedures in the pipeline for the potential user's convenience.

The first step is flatfielding the data. We use a self-flat (i.e., make a flatfield image from the dithered object images) for the  $K_s$  band. An object mask is essential for making good quality flatfields. To do this, we first make “quasi-flatfielded” images by subtracting the adjacent image in time from the image from which we make the object mask. Such a “quasi-flatfielded” image will show a flat sky and objects in the image as positive-negative pairs. Then the residual pattern in the sky of each quasi-flatfielded image is further subtracted by fitting a 3rd-order polynomial surface. This procedure is done independently for each quadrant, because MOIRCS is a 4-channel readout system and any pattern related to the sky variation has discontinuities on quadrant boundaries. After replacing all the negative objects to the sky value, cosmic-ray removal is applied. Each resulting image is then smoothed by Gaussian kernel after a 3 by 3 boxcar median filter has been applied. At this stage, most objects are clearly seen on the smoothed image except very faint ones. Then any area that has a signal larger than the threshold of  $1.25\sigma$

of the original pixel-to-pixel variation are replaced by the “mask” value of 1, with an additional expansion to the area (here we set 5 pixels) for large and bright objects. This quick object mask works well for the usual science exposures. We note that, though we can do the more accurate “second-pass” mask-making procedure after getting the final combined image, the result is almost the same and is not necessary for our data.

After making the object masks, the flat frame is made from the science data itself with the mask applied. For the narrowband data we used dome flat data as we describe further below. After the flatfielding of the NB data, we further subtract the “median sky” image (made with the object masks) which is made from the subset (typically 6 to 10) of the frames before and after the image for the sky subtraction. For the  $K_s$  data this procedure can be skipped because the sky pattern is relatively smooth (by the use of the self flat) and so we can subtract the sky in the next procedure reasonably well. Then we further subtract any residual sky pattern remaining on each image using a 3rd-order polynomial surface fitting for each quadrant.

The resulting flatfielded, sky-subtracted images are then corrected to remove the geometric distortion. The distortion correction database, which is made from observations of astrometric reference fields, is supplied by the Subaru Observatory. We then measure the position of some (typically 20 to 50) bright and compact objects on each image. The object catalogs for the images are further cross-matched with the catalog for the first image in the input list, and from the result we calculate the offset values we use to register the images to the first image. Finally, these images are combined, where we weight by the rms<sup>2</sup> of the input images.

A note on our NB2288 reduction: though we usually use the sky- or self-flats for the reduction, we used the dome flat data for correcting the pixel-to-pixel variation of the NB data. One reason is because the sky counts in the narrow-band filter are generally low compared with that of broad-band data. There is also the possibility that the sky may not be flat in the NB filter. Due to the shift of the observable wavelength window with position (see §2.1.1), the inclusion or exclusion of OH night-sky lines into the window becomes position-dependent. Although there is a small tilt pattern in the dome flats which is caused by the difference of the light path between dome data and object data, we estimate that the variation of the photometry due to the use of dome flat is  $< 3\%$  ( $\sim 0.033$  mag rms) by putting standard stars at several positions across the field of view. This is small compared to our required accuracy.

## 2.3. Photometry and the Catalog

The object detection and the photometry are done using the “double-image” mode of SExtractor 2.5.0 (Bertin & Arnouts 1996). For the detection image, we used the unsmoothed version of the narrowband data; therefore, our catalog is a simple narrowband-based selection. The  $K_s$  image is registered to this, and the seeing size is closely

<sup>2</sup> Available at  
<http://www.naoj.org/staff/ichi/MCSRED/mcsred.html>

<sup>3</sup> <http://iraf.noao.edu/>



matched (6.0 pixel,  $\sim 0.67$  arcsec FWHM) for both images. The object detection is done within the area that is covered by more than 12 frames (75%) for NB2288 as well as 4 frames (67%) for  $K_s$ . For the detection parameters, we use the SExtractor DETECT\_THRESH= 1.75 and DETECT\_MINAREA= 15 with a tophat detection filter (tophat\_4.0\_5x5.conv).

The effective survey area is  $23.6 \text{ arcmin}^2$  which corresponds to a survey volume of  $840.7 \text{ comoving Mpc}^3$  at  $z=2.49$  considering our NB2288 filter bandwidth. We used aperture magnitudes in a diameter of 12 pixels ( $= 1''.4$  or  $11.5 \text{ kpc}$  physical scale) for color measurement, and the MAG\_AUTO values for derivation of the star-formation rates as well as for the stellar mass (§3.5).

The standard star data for FS151 were reduced using the same flatfielding data we used for the object reductions. The instrumental magnitude for FS151 is measured from the individual 4 frames without combining them because the images are slightly defocused. These magnitudes were then averaged to get the final measurement. The instrumental magnitude for each frame is measured by the “growth-curve” method which estimates the convergence magnitude with increasing aperture radius. We used the IRAF APPHOT package.

The  $K_s$ -band magnitude for FS151 is taken from Leggett et al. (2006), with the +1.86 conversion from the Vega to AB-magnitude zeropoint applied (Tokunaga & Vacca 2005). For the NB2288 magnitude, we used the  $K_s$  magnitude and the low-dispersion template spectra of a G-3 dwarf star by Wallace & Hinkle (1997). As a final check, we compared the  $K$ -band magnitude of some unsaturated stars in our catalog to the 2MASS photometry. We find that the magnitudes are well matched at the  $\sim 0.01$  magnitude level with a scatter of  $\sim 0.1$ .

Good photometric error estimates are critical for robust detection of emission-line objects at lower equivalent widths (EWs). In our analysis, we use the following two methods. First, we measure aperture photometry in 1000 random (blank) regions to estimate the  $1\sigma$  level of the sky variation. This yields  $5\text{-}\sigma$  limiting magnitudes in a 12-pixel diameter aperture of 22.3 and 22.9 for  $K_s$  and NB2288, respectively. Second, we estimate the scatter of the  $K_s$ –NB2288 color introduced by the photometric uncertainty. We generated 70000 fake objects with various magnitudes and placed them randomly in blank image regions. After rebinning the sample points in 0.5-magnitude intervals of NB magnitude, the scatter of the output color excess ( $K_s$ –NB2288) is evaluated by measuring the distribution’s 68%( $1\sigma$ ), 95%( $2\sigma$ ), and 99.7%( $3\sigma$ ) levels directly from the simulation. The result of this simulation is used for the estimate of the error on the color excess and on the selection of emitters.

#### 2.4. Supplementary Data Used for Analysis

In the current paper, we use some supplementary datasets (§3.1 & §3.4) and here we briefly describe them, though the full description will be presented elsewhere (I. Tanaka et al. in prep.).

Deep  $J$ -band data were taken using the INGRID camera (Packham et al. 2003) on the William Herschel 4.2-m Telescope (WHT), La Palma. The data were acquired on the photometric nights of UT 2001 Sep 7th and 8th, under good seeing conditions ( $\sim 0.6$  arcsec.). The INGRID camera has a HAWAII HgCdTe  $1024 \times 1024$  array, covering roughly  $4' \times 4'$  field of view ( $= 0''.238/\text{pixel}$ ) on the Cassegrain Focus of the WHT. The total integration time was 4800 s, resulting in a 50% detection completeness limit of 24.4 mag. The data were reduced in the standard manner, similar to the method for MOIRCS data reduction.

Deep  $B$ -band data were taken by the Suprime-cam on Subaru Telescope on the evening of UT 2009 Nov 17th under clear but a marginal seeing condition ( $0''.9 \sim 1''.2$ ). It was taken as part of our Ly $\alpha$  emitter search in this field. The net exposure time was 4500 s, having a limiting magnitude of 26.5 mag ( $5\sigma$ ,  $2''.5$ -diameter aperture). The detail of the data reduction as well as the result of the broadband SED fitting analysis are to be described elsewhere (I. Tanaka et al., in preparation).

MOIRCS MOS spectroscopic observations of this field were done on UT 2006 September 7th (S06B-106: Kodama et al.); i.e., before the acquisition of the narrowband data we describe here. Part of the target selection was done using a similar narrowband survey executed in September 2001 using the INGRID camera on WHT (IT and CP), which should sample the redshift range of  $2.429 < z < 2.489$ . The conditions were clear, with medium ( $\sim 0''.8$ ) seeing conditions. We used the grism with a spectral resolution of  $\sim 1000$  in the  $K$ -band window covering the  $2.0 - 2.4 \mu\text{m}$  spectral range. We used  $0''.8$  wide slits with typical slit lengths of  $\sim 12''$ . Each single exposure was 600 s to 1200 s, depending on the airglow line strength. The standard A-B dither pattern was applied during each exposure. In total 2-hrs of integration were gathered. The MOS data reduction was done in the standard manner using the MCSMDP pipeline (pre-release version) by one of the MOIRCS builders (T. Yoshikawa, private comm.). Details of the MOS reduction procedure by MCSMDP can be found in the recent paper of Yoshikawa et al. (2010).

Finally, the deep  $24 \mu\text{m}$  imaging from *Spitzer*/MIPS were obtained as part of GO program 30240 (PI: A. Stockton). The total exposure time in the central  $2' \times 2'$  of the mosaic is approximately 5 ks. The 171 separate exposures each had a background fitted and then subtracted before a median of the array was determined. This median was then subtracted from each exposure. The exposures were then mosaicked with MOPEX software<sup>4</sup> with standard inputs and resampling the pixels by a factor of 2.

For the current analysis we focus on a subset of the  $24 \mu\text{m}$ -selected sources that have clear IRAC counterparts. The MIPS photometry was extracted using DAOPHOT in IRAF to fit an empirical PSF model to multiple proximate (within  $12''$ ) sources simultaneously. The MIPS source positions were pre-defined using the position of the closest

<sup>4</sup> <http://ssc.spitzer.caltech.edu/dataanalysistools/tools/mopex/>

IRAC counterpart. This photometry method mitigates the effects of source confusion.

### 3. Results

#### 3.1. Emission-line Object Selection

The selection of  $H\alpha$  emitters at  $z \sim 2.5$  is based on excess emission in NB2288 measured via  $K_s$ –NB2288 color. In Figure 2, we show the color-magnitude diagram for all objects detected in our observations. The overall distribution of the detected objects tends to become broader with increasing magnitudes due to the increasing photometric uncertainty. The intrinsic scatter is also seen for the brighter objects. The intrinsic scatter is caused by the variation of the spectral energy distributions (SEDs) in the  $K_s$  band.

The two curves labeled as  $\Sigma = 2$  & 3 in Fig.2 are the conventionally used measures of the significance of narrowband excess (e.g., Geach et al. 2008). It is formulated as below,

$$m_K - m_{NB} = -2.5 \log[1 - \Sigma \delta 10^{-0.4(m_z - m_{NB})}], \quad (1)$$

where  $m_z$  is the zeropoint of the narrowband image and  $\delta$  is the photometric uncertainty. The  $\Sigma = 2$  line generally matches well with our simulated  $\sigma = 2$  curve. As the color scatter becomes large at  $NB2288 > 22$  mag, we limit the selection of emission-line objects to  $\Sigma > 2$  and  $NB2288 \leq 22$  (corresponding to  $7\sigma$  level in 12-pixel aperture). We note that the use of our simulated  $\sigma = 2$  scattering curve does not change the sample. At  $NB2288 > 22$ , there are actually only a few emission-line object candidates with  $\Sigma > 2$ , due to the steep rise of the selection curve.

At the bright end, the intrinsic scatter is relatively large due to the variation of the SEDs in the  $K_s$  band window and the selection of emitters with low-equivalent width may introduce spurious sources. We therefore introduce a further selection criterion using the rest-frame equivalent width,  $EW_o > 50 \text{ \AA}$  at  $z = 2.49$ . This corresponds to the  $K_s$ –NB2288 = 0.55 mag. The limiting  $H\alpha$  line flux of the emitter sample by these criteria is roughly  $7.5 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ . These values are computed using the same formula used in Geach et al. (2008).

We find 12 objects that satisfy the selection criteria set here. Removing one object which is found to be a diffraction spike by visual inspection, the remaining 11 objects are cataloged as robust emission-line objects for further analysis. We note that the brightest object in the emitter sample is 4C 23.56. The observational properties of the emitter sample are summarized in Table 1.

##### 3.1.1. Spectroscopy

Though spectroscopic observations for all the emitter candidates are not obtained yet, four of the 11 emission-line candidates have NIR spectroscopy. These data were taken before the acquisition of the current NB data. The one- and two-dimensional spectra for each target are shown in Figures 3 and 4. HAE#491 is the radio galaxy 4C 23.56 itself.

Due to the relatively low signal-to-noise ratio (SNR) of the spectroscopy, unambiguous identification of  $H\alpha$  via

detection of the nearby [NII] or [SII] lines is only possible for the radio galaxy. The objects HAE#526 and HAE#153 are relatively bright, and we can say this is not [OIII] $\lambda 5007$  emission at  $z = 3.571$  because of the lack of the [OIII] $\lambda 4959$  near  $2.267 \mu\text{m}$  (as well as the lack of  $H\beta$  at  $2.222 \mu\text{m}$ ). Furthermore, the spectra of the latter object seems to have a hint of the broad component which would not appear if it is [OIII]. The lines from these two sources may also be from the Paschen series. If it is the  $Pa\alpha$  ( $Pa\beta$ ) line,  $Pa\beta$  ( $Pa\gamma$ ) will be observed at  $1.56$  ( $1.95$ )  $\mu\text{m}$ . Unfortunately our spectra do not cover wavelengths below  $2 \mu\text{m}$ . We have to make use of other information to exclude the latter possibility (see next section). For the object #354, the  $H\alpha$  signal at  $2.288 \mu\text{m}$  is quite uncertain. However, we see a possible signal at  $2.342 \mu\text{m}$  at the expected position of the [SII] line. Though both lines are quite low SNR, the inferred large [SII]/( $H\alpha$ + [NII]) ratio of  $\sim 1$  would indicate that the object harbours an AGN (Osterbrock 1989). We list the measured redshift values in Table 1 supposing the line is  $H\alpha$ .

The velocity difference among these emitters is relatively small. They lie in the range of  $2.287 - 2.289 \mu\text{m}$  ( $230 \text{ km s}^{-1}$  range), which is in the red edge of the INGRID narrowband filter function we used for pre-selection ( $2.25 - 2.29 \mu\text{m}$ ). If these emission lines are not  $H\alpha$  but from the interlopers unrelated to 4C 23.56, we can expect a more scattered distribution in wavelength range across the INGRID filter function. This is a strong indication that they are physically associated with each other and 4C 23.56.

##### 3.1.2. Broadband Colors of HAEs

Our emission-line survey for  $H\alpha$  at  $z = 2.49$  may contain interlopers at different redshifts. As we showed in the previous section, the spectroscopic detection of a single line remains ambiguous; it could be  $Pa\alpha$  for example. Geach et al. (2008) used color selection as well as photometric redshifts to distinguish interlopers in their  $H\alpha$  emitter sample at  $z = 2.2$ . They estimated that, among their 55 “robust” sample galaxies, a third can be lines other than  $H\alpha$  (12  $Pa\alpha$ , 6  $Pa\beta$ , and one [OIII]). Their result indicates that foreground objects ( $Pa\alpha$  objects at  $z = 0.22$  and  $Pa\beta$  objects at  $z = 0.78$ ) could be the major contaminants of our data.

To make sure that our detected emission-line objects are mainly  $H\alpha$  emitters, we measure the continuum color of our emission-line candidates using our supplementary Subaru  $B$ -band and WHT  $J$ -band data (§2.4). The  $J$ -band data covers the central  $3'8 \times 3'4$  area of the MOIRCS field of view (here we refer to the area as the “INGRID subfield” as shown in Figure 6), while the whole area is covered by  $B$ -band data.

Though our selected emission-line candidates all lie in the INGRID subfield, 4 objects are not seen in our  $B$ -band data. Two more objects are overwhelmed by the PSF of bright nearby stars. We could measure a  $BJK$  color for five objects. Fortunately, all four spectroscopic targets are in this sub-sample. The  $B - J$  versus  $J - K_s$  two-color diagram is shown in Figure 5a. The  $H\alpha$  emitter

candidates are marked by filled boxes (red).

As discussed in detail in the Appendix, the  $B - J$  versus  $J - K_s$  two-color diagram is found to be quite effective for rejecting objects at  $z < 1$ . Figure 5b is the  $BJK$  two-color distribution of galaxies in GOODS-North region from the MOIRCS Deep Survey (Kajisawa et al. 2009; Kajisawa et al. 2010, and references therein). The objects that satisfy  $(J - K_s) > 0.4$   $(B - J) - 0.2$  are empirically shown to have spectroscopic redshift of  $z > 1$  from Fig.5b.

The position of our 5 emission-line objects in Fig.5a is quite favorable; with the exception of one object, they satisfy the  $BJK$  criterion. These 4  $BJK$  emitters comprise our spectroscopic sample. Thus we can conclude that the emission line we detect spectroscopically is indeed H $\alpha$ , not from the Paschen series. This is strong support for the existence of a protocluster at  $z = 2.5$  in the 4C 23.56 field.

One object whose color is below the  $BJK$  line could be an interloper, though we cannot conclude this for certain until its spectra is obtained. One such object can be expected from the field survey as we see in §4.1. During the following discussion we consider that all our selected emission-line objects are HAEs at  $z \sim 2.49$ , though a few of them might be foreground objects.

### 3.2. Spatial Distribution of HAEs

The spatial distribution of our HAE sample on the sky is displayed in Figure 6. The position of 4C 23.56 is marked with the large diamond. The distribution of our selected HAEs (shown with open red circles) is quite tightly clustered at the east to south-eastern side of the radio galaxy.

The biased distribution of HAEs to the eastern (i.e., channel-1) side of 4C 23.56 is not due to the wavelength shift of the filter transmission window or any difference between channels. It is also not due to some unexpected trouble in the channel-2 filter, because 4C 23.56 and the adjacent two emitters are actually detected on the channel-2 side. The inhomogeneous distribution seems to be the reflection of the real distribution of the emitters.

If we suppose that the majority of the protocluster members are strongly redshifted relative to 4C 23.56, the situation could change. As is discussed by §2.1.1, HAEs redshifted by  $>1000$  km/s relative to 4C 23.56 will drop off from our detection window in the outer half of the FOV. In the inner half of the FOV we should detect objects with large velocity difference. Though our three HAEs with spectroscopic redshift have small velocity offset ( $< 230$  km s $^{-1}$ ), further spectroscopy for all the HAEs may uncover a biased velocity distribution.

### 3.3. Offset H $\alpha$ Emission in the “Eastern Clump”

Here we comment on the objects in the area enclosed by the small white box in Fig.6. There is a group of 4 emitters that lies within a 15''-diameter area ( $=300$  kpc comoving), which is also shown in Figure 7 as a zoomed view. We call this area the “Eastern Clump”. The compact size of the HAE group suggests that they might eventually merge into one massive galaxy in the future and become (one of) the brightest cluster member(s).

One notable character of HAEs in the Eastern Clump is that the light distribution of each HAE in the NB2288 image is offset from the  $K$ -band object position, as shown in Fig.7. Especially interesting is that the offset direction of emission-line gas in three HAEs on the west side of the clump in Fig.7 is roughly aligned to the North to North West.

To show these offsets more clearly, we ran SExtractor on each image independently and measured the positional offset of objects detected in both images. The result is shown in Figure 8. In the figure, the 1-, 2-, and 3- $\sigma$  scatter, based on the objects that have similar magnitudes to the HAEs (dots), are shown by circles of dashed lines. The HAEs are marked as filled squares, with the objects in the Eastern Clump shown in Fig.7 overlaid with asterisks. There are two more large-offset ( $> 3\sigma$ ) objects seen in Fig.8. One of them is indeed the object just south of the eastern clump. We confirm that the offset of the emission-line gas is not due to poor astrometric alignment of the two images; all the other objects adjacent to the Eastern Clump show the same scatter shown as dots here.

The  $K$ -band light centroid can be considered the center-of-mass for each galaxy. Therefore, these offsets of the H $\alpha$  emission indicate that the star-formation is occurring at the periphery of the galaxies. This could be due to interactions among the galaxies triggering star-formation (e.g., Sulentic et al. 2001; Cortese et al. 2006), though the apparent alignment of the offset direction is difficult to explain. The apparent offset of the H $\alpha$  emission is roughly  $0''.2 - 0''.4$ , corresponding to a physical size of  $1.6 - 3.2$  kpc at  $z = 2.49$ .

Another possible explanation may be that vast outflows of emission-line gas are being launched from these galaxies. The direction of such outflows could be influenced by the direction that the group is moving towards the center of the cluster potential. If the cluster potential is already filled by dense intracluster gas, the expelled emission-line gas from these galaxies may leave a “wake” behind each galaxy. In the local universe, similarly offset light from emission-line gas is caused by ram-pressure stripping of galaxies by intracluster gas (e.g., Yoshida et al. 2002; Yagi et al. 2007; Sun, Donahue, & Voit 2007). For example, Cortese et al. (2007) discovered a trail of blue emission-line knots behind an object in Abell2667 at  $z = 0.23$ . Our detected offset of the H $\alpha$  emission in the Eastern Clump may be similar.

A detection of extended X-ray emission from 4C 23.56 intracluster gas is the key for falsifying the ram-pressure stripping scenario. The field was observed by *XMM* and reported by Johnson et al. (2007). Though they detected extended soft X-ray emission near the radio galaxy in a  $\sim 30$  ksec EPIC exposures, they concluded that inverse-Compton scattering of CMB photons is the most likely emission mechanism. Indeed the position and direction of the X-ray extension is well aligned with the radio lobe which extends  $\sim 0.5$  Mpc north-east/south-west from the radio galaxy, and not with the direction where the Eastern Clump lies. For now, there is no evidence of thermal X-ray emission from hot intracluster gas.



### 3.4. HAEs and the Spitzer MIPS Data

Spitzer MIPS 24  $\mu\text{m}$  imaging has been used to probe star formation activity at  $z \sim 2$  (e.g., Daddi et al. 2007). There is public, archived deep 24  $\mu\text{m}$  MIPS data for the 4C 23.56 field (§2.4). If the protocluster around 4C 23.56 is characterized by active star-formation, we may also see it as an excess of faint MIPS sources.

In Figure 9 we show our re-processed MIPS 24  $\mu\text{m}$  image. The strong point source at the center of the image is 4C 23.56. We can clearly see an excess of faint MIPS sources around the radio galaxy compared to the outer parts of the image. We also find that 9 objects out of our 11 HAE sample (shown as red open circles) have MIPS counterparts. However, the apparent excess of 24  $\mu\text{m}$  sources does not seem to be due solely to the HAE counterparts. Here we examine the excess of MIPS-selected objects in the 4C 23.56 field. The 24  $\mu\text{m}$  photometry and the inferred star-formation rates for the HAE counterparts are separately discussed in the next section (§3.5).

To estimate the density excess of MIPS sources quantitatively, we have to consider the difference of the exposure coverage across the field. We first exclude any area that has exposure time less than 1/3 of the total integration. We then set the SExtractor object detection threshold to  $2.8\sigma$ . This is higher than usual so that the detection is not affected by the S/N difference due to the variation of exposure time across the field. To check the robustness of the detection threshold further, we reversed the pixel count of the MIPS image and executed the object detection with the same SExtractor parameters. Besides a clear pixel defect, we detected no sources in the inverse image, indicating that the no fake object from the sky variation is included in the current analysis. Furthermore, we exclude all objects with flux density less than  $90 \mu\text{Jy}$ , as they may suffer from the source confusion in crowded regions.

We also introduce an upper flux density cut of  $275 \mu\text{Jy}$  because most of our HAEs are fainter than  $200 \mu\text{Jy}$ . Any object much brighter than  $200 \mu\text{Jy}$  has a high possibility of being in the foreground. The upper flux density cut is chosen to be roughly twice the median 24  $\mu\text{m}$  flux density of our HAEs.

Note that we apply the SExtractor detection directly to the 24  $\mu\text{m}$  data for the source density excess estimate. For the SFR estimate from the MIPS photometry for our HAE sample we use DAOPHOT software as described in §2.4, because it provides a more robust flux estimate of each source.

The resulting density map of the faint ( $90 - 275 \mu\text{Jy}$ ) MIPS sources is overlaid on Fig.9 as contours. The distribution of sources has been smoothed by a Gaussian kernel of  $\sigma = 30''$ . For reference, the position of the HAEs are shown as red open circles. It is quite clear that the MIPS source density excess closely matches the HAE distribution.

To quantify the density excess of MIPS sources further, we define the “excess region” as the  $80''$ -radius region around the peak density contour, and compare the excess within the region with the “outer region” at larger radii.

From the density of faint MIPS sources in the “outer region” we estimate the expected number in the “excess region” as 15.3. On the other hand, we count 39 objects in the “excess region”, yielding a number enhancement of  $2.55^{+0.59}_{-0.48}$ . This corresponds to a  $4.5\text{-}\sigma$  excess by proper Poisson statistics (Gehrels 1986). As a crosscheck, we downloaded the deep MIPS data for GOODS-N/S fields<sup>5</sup> and counted sources in the same manner, yielding the expected count of  $18.2 \pm 0.7$  for an area with the size of the “excess region”. The source counts of deep MIPS data for ELAIS-N1 field by Chary et al. (2004) also shows the expected number of 17.0. Our number is consistent with those surveys, considering the uncertainty by the cosmic variation.

The HAEs with MIPS counterparts (9) can only explain less than half of the total excess count ( $\sim 24$ ). This implies that our shallow  $\text{H}\alpha$  emitter survey may be missing an even larger number of the dust-enshrouded star-forming objects in the excess region. Future follow-up observations using ALMA etc would be quite interesting to understand these hidden dusty sources.

### 3.5. Star-Formation Rate and Stellar Mass of HAEs

In this section we estimate the star-formation rate (SFR) and the stellar mass of our HAE sample. We note, however, that the estimation of the SFRs and the stellar masses for our HAE sample are uncertain and should be treated with caution.

#### 3.5.1. $\text{H}\alpha$ -based Star-Formation Rate of HAEs

It is known that dust absorption lowers the measured SFR from  $\text{H}\alpha$  emission relative to the intrinsic SFR based on radio or FIR emission (e.g., Hopkins et al. 2001; Afonso et al. 2003; Erb et al. 2006; Daddi et al. 2007). The contamination of the  $[\text{N II}]$  6583Å emission line to the narrow-band flux is also an unknown factor (here we set the factor as 0.2). The contribution of AGN to the ionizing radiation also adds uncertainty to the estimate. These uncertainties can be calibrated using future deep spectroscopy of complete HAE samples.

Kennicutt (1998) gives the conversion formula from  $L(\text{H}\alpha)$  to the SFR as follows.

$$\text{SFR} (M_{\odot} \text{ yr}^{-1}) = 7.9 \times 10^{-42} L(\text{H}\alpha) (\text{erg s}^{-1}) \quad (2)$$

Though the conversion assumes the standard Salpeter (1955) Initial Mass Function (IMF), the IMF is known to overestimate the stellar mass. We here use the conversion correction for the Chabrier (2003) IMF by dividing the value by 1.7, for mainly practical reasons, so that our data can be compared to some recent  $\text{H}\alpha$ -based SFR estimates in the literature. Note that the dust extinction should affect the SFR estimate much more than the uncertainty due to  $[\text{N II}]$  contamination. An AGN contribution can also exist for some galaxies; the unusually high derived SFR for 4C 23.56 is clearly due to the AGN contribution to the  $\text{H}\alpha$  flux.

As for the extinction-correction factor, the *BJK* diagram (Fig.5a) is a useful diagnostic for a part of our sam-

<sup>5</sup> available in <http://ssc.spitzer.caltech.edu/spitzermission/observingprograms/legacy>



ple. In the figure we show the  $A_V = 1$  extinction vector using the Calzetti (2001) extinction law. Assuming that the objects are pure disks with dust extinction, the color distribution of the HAEs indicates that the level of the extinction ranges from 0 to  $\sim 1$ . Using the Calzetti (2001) recipe and the conversion of  $A_{H\alpha} = 0.82A_V$ , an extinction correction factor of 1 to 5 is estimated.

For the remaining 6 objects we cannot estimate the extinction. For comparison, the recent study by Onodera et al. (2010) estimated the level of extinction for their *sBzK*-selected H $\alpha$  emission-line objects that seem to have roughly similar stellar masses as our HAEs (§3.5.3). Their extinction estimate is based SED fitting with additional correction as prescribed by Cid Fernandes et al. (2005) and Savaglio et al. (2005). Their Table 4 indicates that the average of the extinction correction factor is 4.9. We applied the correction factors of 4.9 for the objects we could not use the *BJK* diagram. The raw and the extinction-corrected SFR for each HAE is listed in Table 2.

### 3.5.2. Spitzer MIPS Star-Formation Rates

The bolometric infrared luminosity of galaxies correlates very well with other measures of the star-formation rate and may be the most robust measurement of the SFR. The infrared light is emitted by dust-reprocessing of the emitted UV light from young stars and therefore peaks at rest-frame wavelengths between 100 and 300  $\mu\text{m}$ . In lieu of observations at these frequencies, the longest wavelength datapoints need to be extrapolated using assumed SED models. For this radio galaxy field we have used the observed 24  $\mu\text{m}$  imaging to infer the SFRs for the discovered H $\alpha$  emitters.

We used the SED template for Arp 220, a possible low-redshift analogue, to correct the 24  $\mu\text{m}$  to rest-frame 8.0  $\mu\text{m}$  luminosities (Rieke et al. 2009). From these values we estimated the bolometric infrared luminosity using the Chary & Elbaz (2001) and Dale & Helou (2002) SED models, following the prescription by Daddi et al. (2007)<sup>6</sup>. The resulted infrared luminosities from the two SED models are averaged. Finally, we use the Kennicutt (1998)  $L_{\text{IR}}$  to SFR conversion below,

$$\text{SFR} (\text{M}_{\odot} \text{ yr}^{-1}) = 3.0 \times 10^{-44} L_{\text{FIR}} (\text{erg s}^{-1}). \quad (3)$$

Note that the original formula is divided by 1.5 for Chabrier (2003) IMF (Yoshikawa et al. 2010). These conversions and extrapolation result in SFRs which are uncertain by up to factors of a few (Papovich et al. 2007). For our purposes, to estimate the FIR output from likely protocluster galaxies, these uncertainties are tolerable. The *Herschel Space Telescope* has already quantified these uncertainties for other fields (e.g., Rodighiero et al. 2010) and may eventually provide longer wavelength data for these and other protocluster galaxies. The result of the MIPS-based SFR is also listed in Table 2.

The resulting MIPS-based SFRs are significantly (a factor 2-14 with median of 3.6) higher than the SFRs based

on extinction-corrected H $\alpha$ . We will discuss this further in §4.3.

### 3.5.3. Stellar Mass of HAEs

Next we estimate the stellar mass of our HAE emitters. We employ the conversion formula for the stellar mass from the  $K_s$ -band photometry by Daddi et al. (2004). They showed that for galaxies at  $1.4 < z < 2.5$  the stellar mass is reasonably estimated using the formula below,

$$\log(\text{M}_{*}/10^{11} \text{ M}_{\odot}) = -0.4 (K_{\text{Vega}}^{\text{tot}} - K_{11}) + 0.218 [(z - K)_{\text{AB}} - 2.29] \quad (4)$$

where  $K_{11}$  is the  $K$ -band magnitude (Vega scale) of the average  $10^{11} \text{ M}_{\odot}$ -mass objects of their sample ( $=20.14$  for the average redshift of  $\langle z \rangle = 1.9$ ). Considering the difference of the luminosity distance between  $\langle z \rangle = 1.9$  and 2.49, and the average distribution of  $(z - K)_{\text{AB}}$  color of 1 from the Fig.7 of Daddi et al. (2004), we derive the estimate of the stellar mass at  $z = 2.49$  as

$$\log(\text{M}_{*}/10^{11} \text{ M}_{\odot})_{2.49} = -0.4 K_{\text{AB}}^{\text{tot}} + 8.8 \quad (5)$$

where  $K_{\text{AB}}^{\text{tot}}$  is in AB-magnitude scale.

As a check of the validity of the formula, we compare it to the stellar mass estimates based on SED fitting the 12-band photometry for the GOODS-North MOIRCS Deep Survey Catalog (for details of the model fitting, see Kajisawa et al. 2009; Kajisawa et al. 2010). Figure 10 shows the stellar mass versus  $K$ -band magnitude of the cataloged objects in the redshift range  $2.3 < z < 2.6$ , for objects with photometric redshifts (open boxes) and spectroscopic redshifts (crosses). Though the Daddi estimator by equation (4) fits the overall distribution of the photometric redshift sample reasonably well, the slope seems a bit shallower than the data. Therefore we fit all the photo- $z$ -based data in Fig.10 and got the best fit line as

$$\log(\text{M}_{*}/10^{11} \text{ M}_{\odot})_{2.49} = -0.5127 K_{\text{AB}}^{\text{tot}} + 11.312. \quad (6)$$

We use this formula for the conversion from  $K$ -band magnitude to total stellar mass. The rms scatter of the fit is calculated and found to be only 0.2 dex, consistent with the value quoted by Daddi et al. (2004) (but note that the scatter increases to 2.4 dex if we use brighter subsample of  $K_s < 23$ ). The resulting stellar mass for each HAE galaxy is listed in Table 2. We note that the  $K_s$ -band magnitudes for HAEs will be affected by the H $\alpha$  emission. The rest-frame equivalent width of 100  $\text{\AA}$  will affect the mass estimate by 11%. The median value of the offset due to line emission is  $\sim 20\%$ . This is small compared to the mass estimation uncertainty of  $> 0.2$  dex. We also calculate the specific star-formation rate (SSFR) for the HAEs. The results are shown in Table 2.

## 4. Discussion

### 4.1. Comparison to Field Counts

Our discovery of 11 HAE candidates to a limiting flux of  $7.5 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$  in  $23.6 \text{ arcmin}^2$  area should be compared with the number expected in the general field. Geach et al. (2008) provide us with excellent reference

<sup>6</sup> The formula (4) in Daddi et al. (2007) has a possible typo. The constant should be +1.745, not +1.23, to match the description in the text and to reproduce the relation in their Fig.7

data. Their HiZELS survey measured the number density of H $\alpha$  emitters at  $z = 2.23$  over  $0.6 \text{ deg}^2$  using the narrow-band filter (H $_2$ S1,  $\lambda_c = 2.121 \text{ }\mu\text{m}$ ,  $\delta\lambda = 0.021 \text{ }\mu\text{m}$ ) to a line flux limit of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$ . Their survey field is 91.5 times wider than our own and their filter has a similar bandwidth to our NB2288 filter. They detected 180 emitters including non-H $\alpha$  (e.g., Paschen series, [Fe II], [O III]) objects under their selection criteria of *observed* EW of  $50 \text{ }\text{\AA}$  ( $= 14.3 \text{ }\text{\AA}$  in rest frame at  $z = 2.5$ ). As our data is deeper than theirs, we set our detection limit to match theirs. To a NB magnitude of 21.3 mag we find 5 emitters. Next we have to estimate the how many of their 180 emitters would satisfy our threshold of *rest-frame* EW of  $50 \text{ }\text{\AA}$ . In the filter bandwidth they used, the rest-frame EW $_o$  of  $50 \text{ }\text{\AA}$  (at  $z = 2.49$ ) corresponds to a narrowband excess of 0.56. We estimate the number of emitters that satisfy the EW $_o > 50 \text{ }\text{\AA}$  to be about 90 from their Figure 1. Scaling the number to match our field of view, we derive the expected number of the *field* emitters in our field to be  $\sim 1$ . Note that, similar to our sample of HAE candidates, this field count includes interlopers too. Considering our total of 5 emitters, we can say that our field contains 5 times the expected field count.

Recently Geach et al. (2010) have published more generalized HAE counts from their HiZELS and other surveys. Their Table 2 gives the expected field HAE number counts at the limit of  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$  (EW $_o > 10 \text{ }\text{\AA}$ ) to  $z = 2.2$ . Extrapolating their result to  $z = 2.5$  by third-order polynomial and applying our survey size, we get the expected field count in our survey area as  $1.2^{+0.63}_{-0.46}$ . If we again suppose roughly half of them can be observed as HAE in our EW cut of EW $_o > 50 \text{ }\text{\AA}$ , the expected field count could be only 0.6. Thus the result suggests the overdensity even more strongly.

A deep and wide narrowband HAE survey using the new instrument HAWK-I was recently published (Hayes, Schaerer, & Östlin 2010). A field of  $7'5 \times 7'5$  in GOODS South was surveyed using the narrowband filter NB2090 ( $\lambda_c = 2.095 \text{ }\mu\text{m}$ ;  $\delta\lambda = 0.019 \text{ }\mu\text{m}$ ). Their data are extremely deep, with a  $5\sigma$  limiting magnitude of 24.6 AB corresponding to a line flux limit of  $6.8 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ , nearly an order of magnitude deeper than our own data. The inset histogram in their Figure 1 shows that the number of emitters they detected at NB < 22 is 9, under their EW cut of EW $_o > 20 \text{ }\text{\AA}$  (corresponding to the emission-line excess of 0.29). In their filter bandwidth our selection criterion of EW $_o > 50 \text{ }\text{\AA}$  corresponds to an emission-line excess of 0.60 mag. We estimate that 5 of their emitters satisfy our selection criterion of EW $_o > 50 \text{ }\text{\AA}$  from their Figure 1. Considering the difference in survey areas, the expected field count of the NB emitters for our data is estimated to be  $2.5 \pm 1$ . This estimate is consistent with the results from another deep survey, carried out on a smaller field, by Moorwood et al. (2000). Our count of 11 HAEs is therefore  $5.3^{+5.3}_{-2.6}$  times (or  $3\sigma$ ) more than the field count, which is based on the proper treatment of the small number Poisson statistics by Gehrels (1986).

We note that the redshift evolution of the HAE number count suggested by Geach et al. (2010) is quite strong:

extrapolating from redshifts below  $z = 2.2$  to  $z = 2.5$ , the predicted number counts of HAEs with flux larger than  $10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2}$  decreases by 60%, from 8582 to  $\sim 5380 \text{ deg}^{-2} z^{-1}$ . Although the evolution in number density depends on the limiting flux, this correction for the difference in redshifts (from  $z = 2.2$  to 2.5) will also reduce our field count estimate based on Hayes et al. (2010). Our estimate of the overdensity of a factor 5 should therefore be considered conservative.

As we show in Table 2, about half of our HAEs seem to have a stellar mass of  $\gtrsim 10^{11} M_\odot$ . Such massive objects at  $z \sim 2.5$  are relatively rare in the field. For example, if we count such objects using the MODS catalog (§3.5.3), we find only 10 in their  $103\text{-arcmin}^2$  survey area at  $2.3 < z < 2.6$  (Fig.10). By scaling the value to our survey area of  $23.6 \text{ arcmin}^2$  and  $2.47 < z < 2.49$ , we estimate the expected number in our FOV to be 0.15. The excess of such massive galaxies in the 4C 23.56 field is clear. Recalling that these massive galaxies in the 4C 23.56 field are H $\alpha$ -selected, there could be more inactive massive galaxies missed by our search. We are planning to execute multi-wavelength observations of the 4C 23.56 field. The data will help identify such inactive massive galaxies, and will reveal the whole picture of the protocluster.

#### 4.2. Comparison with Other Protoclusters

Although there are several known protoclusters at  $z > 2$ , there is only one other field that has been searched for HAEs. Kurk et al. (2004a) searched for HAEs in their discovered protocluster surrounding PKS 1138-262 at  $z = 2.16$ . Here we compare the number of detected HAEs.

According to their Figure 6 and Table 5, their selection of EW $_o > 50 \text{ }\text{\AA}$  and  $\Sigma > 3$  combined with a magnitude limit of NB $_{\text{Vega}} < 20.2$  would match the selection we use here. There are  $\sim 10$  HAEs meeting these criteria in the PKS 1138-262 field, roughly the same number as the 4C 23.56 field. Their HAE survey field is roughly half of ours (but wider than our “INGRID subfield” where all our HAEs lie). However, if we use the distribution of their Ly $\alpha$  emitters as a proxy for the HAE distribution (their Fig. 10) there would be at most an additional two HAEs. This implies that the richnesses of the 4C 23.56 and the PKS 1138-262 fields are likely comparable.

There are differences in the properties of the HAEs in the PKS 1138-262 and the 4C 23.56 protoclusters, primarily the spatial distribution of the HAEs relative to the radio galaxies. In the PKS 1138-262 system the HAEs are rather homogeneously distributed around the radio galaxy, with some evidence for higher surface density closer to it. In the 4C 23.56 field the radio galaxy lies at the western edge of the HAE distribution. This does not appear to be a result of observational bias as described in §2.1.1 and §3.2.

This may imply a different evolutionary future of the host galaxy of each radio source. For example, the PKS 1138-262 host may evolve into the centrally-dominant cD-like galaxy relatively early while the 4C 23.56 host may remain one of several dominant galaxies until they merge into a single larger galaxy at

the center of the potential. The host of 4C 23.56 is indeed  $\sim 1.5$  magnitudes less luminous in  $K_s$  than the PKS 1138 host ( $K_s = 18.66$  after removing nuclear component and emission-line contribution; Pentericci et al. 1997). Furthermore, the  $K_s = 18.66$  magnitude for the host of PKS 1138 is unusual; its apparent magnitude is as bright as the brightest cluster galaxies at  $z = 1.5$  (Collins et al. 2009). If we use formula (6) in §3.5.3, the stellar mass of the 4C 23.56 and the PKS 1138 hosts are estimated to be  $4.7 \times 10^{11} M_\odot$  and  $5.5 \times 10^{12} M_\odot$ , respectively. For comparison, Seymour et al. (2007) estimated the stellar mass of both host galaxies based on their *Spitzer* multi-band photometry, finding  $3.9 \times 10^{11} M_\odot$  and  $< 1.3 \times 10^{12} M_\odot$ , for 4C 23.56 and PKS 1138 respectively.

Another difference is in the number of HAEs with strong ( $EW_o > 100 \text{ \AA}$ ) emission. In the 4C 23.56 field, 10 of 11 HAEs have rest-frame  $EW_o > 100 \text{ \AA}$ . In the PKS 1138 field the number of objects that satisfy  $EW_o > 100 \text{ \AA}$  and  $NB_{Vega} < 20.2$  is 5. As the equivalent width relates to the H $\alpha$  luminosity, the excess number of strong EW objects in the 4C 23.56 field implies more active star formation occurring in that protocluster. We will address this quite interesting topic more in the next section.

Recently, Hayashi et al. (2010) reported an excess of [O II] emitters in the field of the distant X-ray cluster XMMXCS J2215.9-1738 at  $z = 1.46$ . The distribution of [O II] emitters are prevalent from the outskirts to the core region of the X-ray cluster. They showed that the fraction of the [O II] emitters among all cluster members as well as the level of the star-formation are shown to be both nearly constant as a function of clustercentric radius (see also, Tran et al. 2010).

Their Figs.5-7 indicate that the distribution of the emitters well traces the overall distribution of all cluster members. The distribution of H $\alpha$  emitters in Kurk et al. (2004a) also closely matches the photo- $z$ -selected galaxy distribution as shown by their Fig.6 (Tanaka et al. 2010). We note, however, that the case of the cluster XMMXCS J2215.9-1738 may not be the representative as a cluster at  $z \sim 1.5$ . Hilton et al. (2010) has shown that the system has a possible bimodal kinematic structure, with the position of the cluster BCG significantly offset from the center of the thermal hot X-ray gas. And also, the NB HAE survey for another cluster XMMU J2235.3-2557 at  $z \sim 1.4$  by Bauer et al. (2010) has show that the HAE distribution just avoids the central  $\sim 200$  kpc of the cluster core (see also Lidman et al. 2008; Strazzullo et al. 2010). Although we are still not sure about whether the HAE can trace the overall distribution of cluster galaxies from the core to the outskirts at  $z > 2$  universe, both studies could consider the supporting case that the distribution of HAEs in the 4C 23.56 protocluster also traces the underlying galaxy distribution. Future analysis based on the broadband imaging will enable us to address the issue.

Hayashi et al. (2010) contrast the activity of galaxies in the cluster core region by comparing to similar work for a  $z = 0.8$  cluster by Koyama et al. (2010). At even

lower redshift, Kodama et al. (2004) studied the HAE distribution for Cl 0024+1652 at  $z = 0.4$ , showing quite strong suppression of activity in the cluster core region. It is quite interesting to connect the results by these works from  $z = 0.4$  (Kodama et al. 2004),  $z = 0.8$  (Koyama et al. 2010),  $z = 1.4 - 1.5$  (Hayashi et al. 2010; Bauer et al. 2010),  $z = 2.1 - 2.2$  (Kurk et al. 2004a; Gobat et al. 2010), to our  $z = 2.5$  studies as a snapshot of the cluster evolution in each epoch (Tadaki et al. 2010). The implied rise of star-formation activity in the cluster core with redshift and its variety suggests that, while it may indicate that we are gradually approaching the era of galaxy formation for members in the cluster core (Kodama & Bower 2001), it is important to evaluate the physical state (richness, dynamics, mass etc.) for each protoclusters in detail to draw a general pictures of early history of galaxy evolution in cluster core. The detailed studies of larger samples are clearly needed to corroborate this apparent evolutionary trend.

#### 4.3. Star-Formation Activity of HAEs

In §3.5 we calculate the SFR and the stellar mass of our HAE sample. Although we admit that the estimate of these values is rather crude, here we try to characterize the star-formation of the HAEs in 4C 23.56 field by comparing them to the other studies. As is discussed in §3.5, the main uncertainty for H $\alpha$ -based SFR is in the estimate of the dust extinction, as indicated by the MIPS detection. The unknown fraction of the AGN contribution to our HAE sample (it can be non-negligible: see Lehmer et al. 2009; Digby-North et al. 2010; Lemaux et al. 2010) is also an issue we need to address by the future observation.

In Figure 11a, we plot the estimated star-formation rate of our HAEs against their stellar mass. The minimum SFR in our sample is  $\sim 15 M_\odot \text{ yr}^{-1}$ , with a median value of 45 (182 for extinction corrected, 746 for MIPS)  $M_\odot \text{ yr}^{-1}$ . In the figure, the brightest object with the highest SFR, is 4C 23.56 (note that the mass estimate for this object is probably overestimated, not only due to the inclusion of AGN but also due to the effect of the poor sky subtraction: see §2.1). The line on the figure is the average mass-SFR relation for  $z \sim 2$  BzK-selected galaxies with MIPS detections by Daddi et al. (2007). The SFRs based on both indicators show larger values than expected from the  $z \sim 2$  mass-SFR relation by Daddi et al. (2007). Especially, the MIPS-based SFR shows large excess. This trend seems real, because our MIPS-based SFR is derived from the same equation as used by Daddi et al. (2007). This may indicate that some galaxies in the protocluster 4C 23.56 have higher star-formation activity than the field counterparts with similar mass.

The observed large excess of the  $24\mu\text{m}$ -based SFR relative to H $\alpha$ -based one may indicate that the star-formation activity in our HAE sample is largely (or selectively) hidden by dust. At lower redshifts such hidden star-formation activity revealed by infrared observation is often reported (e.g., Valdés et al. 2005; Geach et al. 2006; García-Marín et al. 2009; Garn et al. 2010). On the other hand, the recent calibration of the MIPS  $24\mu\text{m}$ -based SFR for star-



forming galaxies at  $z \sim 2$  by the *Herschel* PACS observation has revealed that the SFRs from  $24\mu\text{m}$  alone could be overestimated by a factor of  $\sim 2 - 7.5$  (Nordon et al. 2010; Rodighiero et al. 2010). Our observed trend can be due to the large systematics in the estimate of the total infrared luminosity from the rest-frame  $8\mu\text{m}$  photometry. For now we do not know whether the HAEs in the 4C 23.56 field harbours the star-formation significantly hidden by dust. The direct far-infrared observation by ALMA will address the issue.

In Fig.11b we show the  $L(\text{H}\alpha)$ -based specific star-formation rate (SSFR) as a function of stellar mass. For comparison, we plot the SSFR values for field  $\text{H}\alpha$  emitters from Förster-Schreiber et al. (2009) and Onodera et al. (2010). The Förster-Schreiber et al. (2009) points include two types of the extinction correction, which is shown as connected points with vertical lines in the figure, while Onodera et al. (2010) used the correction (described above) which could be considered the middle between the two extreme cases from Förster-Schreiber et al. (2009).

The measured relation between SSFR and mass deviates clearly from that observed in similar studies of low redshift clusters. For example, Patel et al. (2009) measured the SFR and SSFR of a rich cluster at  $z \sim 0.8$  from MIPS-based photometry, and confirmed that the SSFR of massive ( $\log M/M_\odot > 10.8$ ) galaxies in high density regions is as low as for field objects with similar mass in local universe. Also at  $z > 1$ , the SSFR estimate of the member galaxies in the irregular cluster CL 0332-2742 at  $z \sim 1.6$  by Kurk et al. (2009) show that significant member galaxies have much lower SSFR than field counterparts at similar redshifts, despite that the system is quite irregular and far from a virialized structure. Recently, Tanaka et al. (2010) studied galaxies in the protocluster around PKS 1138 at  $z = 2.16$  and showed a similar trend of the suppressed star-formation activity in the cluster compared to the field environment (see also, Zirm et al. 2008). These studies may indicate that the star-formation activity in cluster environment is suppressed up to  $z \sim 2$ .

To avoid the uncertainties introduced by estimating the SFR and the stellar mass, we also compare the star-formation rates directly using the observed values. In Figure 12 we show the observed  $K$ -band magnitude versus the  $\text{H}\alpha$  luminosity ( $L(\text{H}\alpha)$ ) of the same dataset as shown in Fig.11b, but limited to objects at  $z > 2$ . Here the data from Kurk et al. (2004a) is also added for comparison. Within the uncertainties, all HAEs in the 4C 23.56 field are clearly as luminous as the brightest  $\text{H}\alpha$  sample in the general field results, again suggesting an excess of active star formation in the 4C 23.56 field.

Perhaps more interestingly, in Fig.12 we compare the HAEs in the 4C 23.56 field with the PKS 1138 field data. The majority of HAEs in the PKS 1138 protocluster have lower  $\text{H}\alpha$  luminosities than both field objects and the HAEs in 4C 23.56 despite being within a similar  $K$  magnitude range. It indicates that the HAEs in the PKS 1138 field show suppressed star formation. This is actually the same conclusion reached by Tanaka et al. (2010), but their work is based on broadband SED fitting, i.e., from an in-

dependent measure.

The cosmic time between the two protoclusters is only 0.4 Gyr. This is actually long enough to discern changes of the star-formation activity if their star-formation timescale is short. Tanaka et al. (2010) suggest that the star-formation timescale for galaxies in the PKS 1138 field is much shorter than for galaxies with similar mass in the GOODS fields. Their Figure 13 demonstrates the rapid decline of the SFR in the protocluster compared to that of GOODS, and shows that the time difference of 0.4 Gyr is long enough to observe the decline. For example, if we suppose that these star-forming galaxies in protoclusters are formed at  $z \sim 3$ , the increase of the star-formation rate between  $z = 2.16$  to 2.5 (0.4 Gyr) is close to an order of magnitude.

This argument could of course be too simplified, as it assumes that any galaxies in the two clusters have almost the same formation period around  $z \sim 3$ . The actual formation of massive galaxies in the cluster environment could be much more complicated, and it could be coupled with the fundamental cluster properties (mass, merger history, dynamical state, and so on). As we already discussed, the difference of the star-formation activity between the two protoclusters might be related to the different inherent cluster characteristics. The PKS 1138 host is exceptionally massive and already lies near the apparent center of the galaxy distribution (Kurk et al. 2004a; Seymour et al. 2007), while the 4C 23.56 host lies at the edge of the HAE (and presumably overall galaxy) distribution (§3.2 & §4.2). The soft X-ray emission from the hot intracluster gas is (marginally) detected in PKS 1138 (Carilli et al. 2002), while for 4C 23.56 it is not (Johnson et al. 2007). The future of the PKS 1138 protocluster may well be an X-ray bright, well evolved rich cluster with a massive cD galaxy lying in the center, while that of 4C 23.56 might be an X-ray faint cluster with a few dominant giant ellipticals near the center of the potential, such as the Coma or Virgo clusters.

## 5. Summary

We have discovered a significant excess of near-infrared selected emission-line objects around the radio galaxy 4C 23.56 at  $z = 2.486$ . We detect 11 robust emission-line galaxies with a line flux greater than  $7.5 \times 10^{-17} \text{erg s}^{-1} \text{cm}^{-2}$  in our MOIRCS imaging observation of the  $4' \times 7'$  field around the radio galaxy. We show that this is a significantly higher (5 times more) detection rate compared to other near-infrared emission-line surveys in non-radio-galaxy fields. Three of the emission-line galaxies are found to have a spectroscopic redshift of  $z = 2.49$ . We therefore conclude there is a protocluster physically associated with 4C 23.56.

The distribution of HAEs on the sky is confined to a relatively small area ( $\sim 1.2 \text{Mpc}$ -radius region) near the radio galaxy, with 4C 23.56 placed at the western edge. We show that this is not due to observational bias, and conclude that the protocluster is physically associated with the radio galaxy. The offset position of the radio galaxy

is interesting, because the hosts of powerful radio galaxies are usually very massive and often assumed to be the dominant object (e.g., cD galaxy) in the center of the cluster. We also find that there is an interesting compact group of four HAEs, all the members of which show offset H $\alpha$  emission relative to the  $K_s$ -band light position. This may indicate strong star-formation on the edge of each galaxy which is occurring due to strong interactions with the group. A more exotic interpretation might be that there is ram-pressure stripping of the galactic-scale H $\alpha$ -emitting gas by the hot intracluster gas. Further follow-up is necessary to understand the phenomenon.

Our analysis of deep *Spitzer* MIPS 24  $\mu$ m data for 4C 23.56 shows that all but two of the HAEs have 24  $\mu$ m counterparts. The inferred star-formation rate (SFR) based on the MIPS 24  $\mu$ m photometry is quite high (median of  $> 700 M_{\odot} \text{ yr}^{-1}$ ), indicating that intense starburst activity is going on in each HAE. The MIPS-based SFR is on average 3.5 times higher than the value inferred from the observed H $\alpha$  luminosity. The strong effect of “selective” dust extinction is evident for these galaxies, implying the metal-rich and evolved populations for HAEs, as suggested by Kurk et al. (2004a).

The analysis of the spatial distribution of the MIPS sources has also revealed that there exists a strong excess around 4C 23.56. Notably, the distribution of sources that have 24  $\mu$ m fluxes as faint as the HAEs (90 – 275  $\mu$ Jy) shows the same offset distribution relative to the radio galaxy, and the position of the density peak nicely coincides with the HAE distribution. This is another indication that the strong star-formation activity is going on in the protocluster.

The comparison of the star-formation rates of HAEs in the protocluster 4C 23.56 at  $z = 2.48$  to those in another protocluster PKS 1138-262 at  $z = 2.16$  and to those in the general field at  $z \sim 2$  shows that: (1) HAEs in the 4C 23.56 field are as active as those in the general field, and that (2) HAEs in the protocluster PKS 1138-262 show a clear *deficit* of objects with high H $\alpha$  luminosities. The clear difference in star forming activities between the two protoclusters at similar redshifts may indicate that we are stepping into the epoch of major star formation in massive cluster galaxies. Given the fact that there is a small but significant time difference of 0.4 Gyr between the two redshifts, such variation in star formation activity may be interpreted as a time evolution where star formation activities to form massive galaxies can decay rather sharply during this epoch ( $2 < z < 3$ ) in proto-clusters (e.g., Kodama et al. 2007; Tanaka et al. 2010). Alternatively, we may be just seeing the intrinsic scatter of cluster properties at this high redshift where the onset of cluster formation and the subsequent evolution depend on the mass of the systems and/or structures in the surrounding environment. Such dependence is actually inferred in lower-redshift clusters at  $z < 1$  (e.g., Urquhart et al. 2010). However, we do not know yet the global properties of these protoclusters, such as total number of member galaxies, cluster masses, and the dynamical states, nor do we yet have a statistical sample of protoclusters at this interesting redshift regime to

disentangle these two effects. We are desperate to obtain such information by extensive near-infrared spectroscopy and to increase the sample of proto-clusters at  $z > 2$ .

We thank the anonymous referee for comments that has lead to the improvement of the paper. We thank the MOIRCS Deep Survey Team for providing us with the MODS catalog. Also, we would like to thank Dr. Sune Toft and Dr. Masa Tanaka, and many others for useful comments and suggestions. We also thank Dr. Ryuji Suzuki for giving simulation data on the shift of the central wavelength of NB filters. This work was supported by the JSPS Institutional Program for Young Researcher Overseas Visits. YT and TK thank for support via Grant-in-Aid for Scientific Research (KAKENHI: Nos. 19340046 and 21340045, respectively) by the Ministry of Education, Culture, Sports, Science and Technology in Japan. JK thanks the DFG for support via German-Israeli Project Cooperation grant STE1869/1-1.GE625/15-1. IT would like to thank the all ESO staff for the hospitality during the stay in ESO Garching. IT would also like to thank all the staffs and members of the DARK Cosmology Center, MSSL University College London, and Durham University for their warm hospitality during my stay related to this study. This work is based on data collected at Subaru Telescope, which is operated by the National Astronomical Observatory of Japan. This is also based in part on observations made with the *Spitzer* Space Telescope which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA, as well as the data collected by the 4.2m William Herschel Telescope which is operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

## Appendix 1. The *BJK* Color Selection for Rejecting Objects at $z < 1$

In order to reject the possibility that our single line detection by NB filter or the spectra is mainly from the low-redshift Paschen series interlopers, we introduce the broadband color selection by the  $B - J$  versus  $J - K_s$  two-color diagram. Here we show its effectiveness using the large photometric/spectroscopic catalog for the GOODS-North region.

We make use of the MOIRCS Deep Survey Catalog which contains 13-band photometric data as well as the extensive compilation of spectroscopic redshift data (Kajisawa private com.; Kajisawa et al. 2010). In Figure 5a, we plot the color of all galaxies with  $K_s < 23$  in GOODS-N (gray crosses). The galaxies at  $0.2 < z < 0.25$  and  $0.75 < z < 0.8$  (spectroscopically measured), each of which includes the  $\text{Pa}\alpha$  and  $\text{Pa}\beta$  interlopers, are shown by blue filled triangles, while the objects at  $2.3 < z < 2.6$ , which include the 4C 23.56 redshift, are shown by red filled circles. There is a clear line between the two samples. Here we define it as,

$$(J - K_s) = 0.4 (B - J) - 0.2, \quad (\text{A1})$$

which is shown as the magenta dotted line in the figure. The insetted diagram in Fig.5b shows the fraction of galaxies whose  $J - K_s$  color is redder than the line defined here (“*BJK*” galaxies) as a function of redshifts (spectroscopic redshift sample only). Almost no objects at  $z < 1.2$  can be *BJK* galaxies, while more than 80% of objects at  $z \sim 2.5$  satisfy the criterion.

We also calculate the behavior of galaxy colors in the *BJK* plane using the population synthesis SED models by Kodama & Arimoto (1997) overplotted in Fig.5a. In the figure, each model track indicates, from bluer to redder, the “pure disk” model with continuous SFR (bluest), “50% disk + 50% bulge”, “20% disk + 80% bulge”, and “pure bulge” model with the formation redshift of 5 (reddest, and mostly out of the figure). In each model, symbols are put from  $z = 0.1$  to 4, with the interval of 0.5 from  $z = 0.5$  (each line starts from  $z = 0.1$  with a symbol, and ends at 4.5 without symbol). A part of each model track has the region with cyan-colored solid line, which indicates the region at  $2 < z < 3.8$ . The extinction vector for  $A_V = 1.0$  based on Calzetti (2001) is also plotted.

The behavior of the model tracks also confirms that the our proposed dividing line for *BJK* galaxies works rejecting objects at  $z < 1$ . The model tracks at  $2.3 < z < 2.6$  (the same redshift range as red filled circles in Fig.5b) tend to go a bit further from the *BJK* line than the actual distribution of objects at the same redshift range in Fig.5b. This may be explained, though, by a difference in the metallicity between the modeled and observed galaxies. More advanced modeling is beyond the scope of this paper, however.

## Appendix 2. Objects Identified in Previous Studies

As is described in §1, the 4C 23.56 field was also observed previously by KC97, Kajisawa et al. (2006b), and Stockton et al. (2004). Here we briefly comments on the objects detected in each study.

### A.2.1. Objects in KC97

KC97 studied the  $1.25 \text{ arcmin}^2$  region around the radio galaxy using deep *U'VIR* imaging data from 2.2- & 3.6-m telescopes. They also searched for the  $\text{Ly}\alpha$  emitters using the narrowband filter at  $4229\text{\AA}$  on the UH 2.2-m telescope. They cataloged 9 possible  $\text{Ly}\alpha$ -emitting companions and 15 EROs selected by  $I - K_s$  colors around the radio galaxy.

We first looked for the counterparts of their  $\text{Ly}\alpha$  emitters in our  $K_s$  image. However, we can only identify one object (ID#110 in their catalog). The  $K_s - \text{NB}$  color of the object is 0.14, so no clear sign of NB excess is seen in Figure 13 (red filled box).

Next we identify their EROs. Of their 15 objects, 6 objects are not seen at all in our  $K_s$  nor NB images, and 5 objects are quite faint and subsequently not in our SExtractor catalog. Note that their data is extremely deep, with a  $3\sigma$  limiting magnitude of  $K = 24.3 \text{ mag}$ , while our data only reaches  $23.5 \text{ mag}$ . Among the 4 remaining objects, only one object (their ID#70) shows a sign of emission-line excess (green open circles in Fig.13). However, the object is just below our HAE selection criteria ( $\text{NB} < 22 \text{ mag}$  and  $\Sigma > 2$ ) and therefore not present among our HAE candidates.

We comment on the failure of the HAE counterpart detection for their  $\text{Ly}\alpha$  emitters in detail. We can calculate the expected  $\text{H}\alpha$  flux from their cataloged equivalent width, assuming the  $\text{Ly}\alpha/\text{H}\alpha$  ratio of 8.7 for the Case B recombination (Osterbrock 1989). We find that the expected  $\text{H}\alpha$  flux are mostly several  $\times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ , close to our detection limit. However, this should be seen as a lower limit, as the  $\text{Ly}\alpha/\text{H}\alpha$  can be much smaller than 8.7 due to the more effective dust absorption by the resonant scattering nature of the  $\text{Ly}\alpha$  line (Kurk et al. 2004a; Hayes et al. 2010). Thus at least some, if not all, of their 9 LAEs could be detected. We note that the cataloged  $K$ -band magnitudes of  $\text{Ly}\alpha$  emitters are all quite faint (5 are upper limits with  $K_{\text{AB}} > 24.35$ , and only two are within our magnitude limit of  $K_{\text{AB}} > 23.5$ ). The non-detection by our  $K$  image could be reasonable. However they are also totally absent even in our NB2288 image.

Kurk et al. (2004a) also did not detect  $\text{H}\alpha$  emission for their sample of  $\text{Ly}\alpha$  emitters in the PKS 1138 protocluster. For two of their three brightest  $\text{Ly}\alpha$  emitters, for which  $\text{H}\alpha$  emission could be detectable, faint emission was seen, but not strong enough to pass the HAE selection criteria. The range of the upper limit of the  $\text{Ly}\alpha/\text{H}\alpha$  ratio for their  $\text{H}\alpha$  emitters are  $0.03 - 1.06$ , significantly lower than the Case-B assumption (see also, Hayes et al. 2010).

There are two possibilities to explain this. One possibility is that the redshift of the  $\text{Ly}\alpha$  emitters are out of our NB2288 narrowband window. The filter used by KC97



has a central wavelength of 4229Å with a filter width of 70Å. The Ly $\alpha$  emission that can enter into the window should be in the range  $2.45 < z < 2.50$ , and the corresponding H $\alpha$  line will come to  $2.261 \mu\text{m} - 2.294 \mu\text{m}$ . As is shown in Fig.1, almost half of the shorter wavelength range ( $2.261 \mu\text{m} - 2.28 \mu\text{m}$ ) is out of our survey window. However, if the Ly $\alpha$  emitters are all such objects, they would not be associated with, but slightly foreground to 4C 23.56.

Another explanation is a large error in the Ly $\alpha$  EW estimate, or spurious detections. Due to the lack of *B*-band data which samples the continuum just redwards of Ly $\alpha$ , they had to use a crude UV continuum estimate. If the EW of their Ly $\alpha$  emitters are much smaller than their quoted values, our non-detection of H $\alpha$  counterparts could be understood. An independent Ly $\alpha$  emitter search is necessary to check the reliability of their selection of NB excess objects. We are in the progress of carrying out such a Ly $\alpha$  imaging survey for this field. Its results will be published elsewhere (Tanaka I. et al., in preparation).

#### A.2.2. *Objects in Kajisawa et al. (2006b)*

Kajisawa et al. (2006b) took the *JHK* images of the 4C 23.56 field and found a significant excess of “*JHK*” objects (see also, Kodama et al. 2007). We checked their H $\alpha$  excess in our catalog. Though we could not identify their *b-JHK* objects, several Distant Red Galaxies (DRGs) were identified. The color excess of their DRGs is shown in Fig.13 (blue filled circles). Excluding 4C 23.56, two objects are identified in our HAE sample. Another two objects are identified as KC97’s EROs. One of them shows emission-line excess (#70 in KC97). Two other objects do not show detectable emission-line excess.

#### A.2.3. *An Object in Stockton et al. (2004)*

Stockton et al. (2004) reported an object in 4C 23.56 having very red color and a disk-like morphology. The SED fitting result indicates the strong preference for the single-burst model with fairly old age of  $\sim 2\text{Gyr}$  at the redshift of the radio galaxy. The result suggests that the galaxy should be inactive Stockton & McGrath (2007), and no H $\alpha$  emission is expected.

Their object is actually identified as an ERO in KC97 which is a DRG in Kajisawa et al. (2006b). In Fig.13 we show the color excess of the object (magenta open triangle). As expected, the object does not show signs of excess emission from H $\alpha$ .


## References

- Afonso, J., Hopkins, A., Mobasher, B., & Almeida, C. 2003, *ApJ*, 597, 269  
 Balogh, M. L., Morris, S. L., Yee, H. K. C., Carlberg, R. G., & Ellingson, E. 1997, *ApJL*, 488, L75  
 Bauer, A. E., Grutzbauch, R., Jorgensen, I., Varela, J., & Bergmann, M. 2010, *MNRAS*, in press (arXiv:1010.1238)  
 Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393  
 Blanton, M. R., & Moustakas, J. 2009, *ARA&A*, 47, 159  
 Bower, R. G., Lucey, J. R., & Ellis, R. S. 1992, *MNRAS*, 254, 601  
 Bower, R. G., Kodama, T., & Terlevich, A. 1998, *MNRAS*, 299, 1193  
 Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 100  
 Bundy, K., et al. 2006, *ApJ*, 651, 120  
 Butcher, H., & Oemler, A., Jr. 1978, *ApJ*, 219, 18  
 Butcher, H., & Oemler, A., Jr. 1984, *ApJ*, 285, 426  
 Calzetti, D. 2001, *PASP*, 113, 1449  
 Carilli, C. L., Harris, D. E., Pentericci, L., Röttgering, H. J. A., Miley, G. K., Kurk, J. D., & van Breugel, W. 2002, *ApJ*, 567, 781  
 Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562  
 Clemens, M. S., Bressan, A., Nikolic, B., Alexander, P., Annibali, F., & Rampazzo, R. 2006, *MNRAS*, 370, 702  
 Clemens, M. S., Bressan, A., Nikolic, B., & Rampazzo, R. 2009, *MNRAS*, 392, L35  
 Cortese, L., et al. 2007, *MNRAS*, 376, 157  
 Cortese, L., Gavazzi, G., Boselli, A., Franzetti, P., Kennicutt, R. C., O’Neil, K., & Sakai, S. 2006, *A&A*, 453, 847  
 Couch, W. J., Ellis, R. S., Sharples, R. M., & Smail, I. 1994, *ApJ*, 430, 121  
 Chabrier, G. 2003, *PASP*, 115, 763  
 Chary, R., et al. 2004, *ApJS*, 154, 80  
 Cid Fernandes, R., Mateus, A., Sodré, L., Stasińska, G., & Gomes, J. M. 2005, *MNRAS*, 358, 363  
 Collins, C. A., et al. 2009, *Nature*, 458, 603  
 Daddi, E., Cimatti, A., Renzini, A., Fontana, A., Mignoli, M., Pozzetti, L., Tozzi, P., & Zamorani, G. 2004, *ApJ*, 617, 746  
 Daddi, E., et al. 2007, *ApJ*, 670, 156  
 Dale, D. A., & Helou, G. 2002, *ApJ*, 576, 159  
 De Breuck, C., van Breugel, W., Stanford, S. A., Röttgering, H., Miley, G., & Stern, D. 2002, *AJ*, 123, 637  
 De Breuck, C., et al. 2004, *A&A*, 424, 1  
 De Lucia, G., et al. 2004, *ApJL*, 610, L77  
 De Lucia, G., et al. 2007, *MNRAS*, 374, 809  
 Digby-North, J. A., et al. 2010, *MNRAS*, 407, 846  
 Doherty, M., et al. 2010, *A&A*, 509, 83  
 Dijkstra, M., & Loeb, A. 2009, *MNRAS*, 400, 1109  
 Dressler, A. 1980, *ApJ*, 236, 351  
 Dressler, A., Oemler, A., Jr., Butcher, H. R., & Gunn, J. E. 1994, *ApJ*, 430, 107  
 Dressler, A., et al. 1997, *ApJ*, 490, 577  
 Eke, V. R., Navarro, J. F., & Frenk, C. S. 1998, *ApJ*, 503, 569  
 Elbaz, D., et al. 2007, *A&A*, 468, 33  
 Elbaz, D., Cesarsky, C. J., Chanial, P., Aussel, H., Franceschini, A., Fadda, D., & Chary, R. R. 2002, *A&A*, 384, 848  
 Ellingson, E., Lin, H., Yee, H. K. C., & Carlberg, R. G. 2001, *ApJ*, 547, 609  
 Ellis, R. S., Smail, I., Dressler, A., Couch, W. J., Oemler, A., Jr., Butcher, H., & Sharples, R. M. 1997, *ApJ*, 483, 582  
 Ellis, S. C., Jones, L. R., Donovan, D., Ebeling, H., & Khosroshahi, H. G. 2006, *MNRAS*, 368, 769  
 Erb, D. K., Steidel, C. C., Shapley, A. E., Pettini, M., Reddy, N. A., & Adelberger, K. L. 2006, *ApJ*, 647, 128  
 Förster Schreiber, N. M., et al. 2009, *ApJ*, 706, 1364  
 Franx, M., et al. 2003, *ApJL*, 587, L79  
 García-Marín, M., Colina, L., & Arribas, S. 2009, *A&A*, 505, 1017  
 Garn, T., et al. 2010, *MNRAS*, 402, 2017  
 Geach, J. E., et al. 2006, *ApJ*, 649, 661  
 Geach, J. E., Smail, I., Best, P. N., Kurk, J., Casali, M., Ivison, R. J., & Coppin, K. 2008, *MNRAS*, 388, 1473  
 Geach, J. E., et al. 2010, *MNRAS*, 402, 1330  
 Gehrels, N. 1986, *ApJ*, 303, 336


- Gilbank, D. G., Yee, H. K. C., Ellingson, E., Gladders, M. D., Loh, Y.-S., Barrientos, L. F., & Barkhouse, W. A. 2008, *ApJ*, 673, 742
- Gobat, R., et al. 2010, *aap*, submitted (arXiv:1011.1837)
- Gómez, P. L., et al. 2003, *ApJ*, 584, 210
- Hashimoto, Y., Oemler, A., Jr., Lin, H., & Tucker, D. L. 1998, *ApJ*, 499, 589
- Hayashi, M., Kodama, T., Koyama, Y., Tanaka, I., Shimasaku, K., & Okamura, S. 2010, *MNRAS*, 402, 1980
- Hayes, M., Schaerer, D., & Östlin, G. 2010, *A&A*, 509, L5
- Hayes, M., et al. 2010, *Nature*, 464, 562
- Hilton, M., et al. 2009, *ApJ*, 697, 436
- Hilton, M., et al. 2010, *ApJ*, 718, 133
- Holden, B. P., Stanford, S. A., Eisenhardt, P., & Dickinson, M. 2004, *AJ*, 127, 2484
- Hopkins, A. M., Connolly, A. J., Haarsma, D. B., & Cram, L. E. 2001, *AJ*, 122, 288
- Ichikawa, T., et al. 2006, *Proc. SPIE*, 6269,
- Ideue, Y., et al. 2009, *ApJ*, 700, 971
- Johnson, O., Almaini, O., Best, P. N., & Dunlop, J. 2007, *MNRAS*, 376, 151
- Kajisawa, M., et al. 2000, *PASJ*, 52, 61
- Kajisawa, M., Kodama, T., Tanaka, I., Yamada, T., & Bower, R. 2006, *MNRAS*, 371, 577
- Kajisawa, M., et al. 2006, *PASJ*, 58, 951
- Kajisawa, M., et al. 2009, *ApJ*, 702, 1393
- Kajisawa, M., et al. 2010, *ApJ*, submitted.
- Keel, W. C., Cohen, S. H., Windhorst, R. A., & Waddington, I. 1999, *AJ*, 118, 2547
- Kennicutt, R. C., Jr. 1998, *ARA&A*, 36, 189
- Knopp, G. P., & Chambers, K. C. 1997, *ApJS*, 109, 367
- Kodama, T., & Arimoto, N. 1997, *A&A*, 320, 41
- Kodama, T., Arimoto, N., Barger, A. J., & Arag'ón-Salamanca, A. 1998, *A&A*, 334, 99
- Kodama, T., & Bower, R. G. 2001, *MNRAS*, 321, 18
- Kodama, T., et al. 2004, *MNRAS*, 350, 1005
- Kodama, T., Tanaka, I., Kajisawa, M., Kurk, J., Venemans, B., De Breuck, C., Vernet, J., & Lidman, C. 2007, *MNRAS*, 377, 1717
- Koyama, Y., Kodama, T., Shimasaku, K., Hayashi, M., Okamura, S., Tanaka, I., & Tokoku, C. 2010, *MNRAS*, 403, 1611
- Kurk, J. D., et al. 2000, *A&A*, 358, L1
- Kurk, J. D., Pentericci, L., Röttgering, H. J. A., & Miley, G. K. 2004, *A&A*, 428, 793
- Kurk, J. D., Pentericci, L., Overzier, R. A., Röttgering, H. J. A., & Miley, G. K. 2004, *A&A*, 428, 817
- Kurk, J., et al. 2009, *A&A*, 504, 331
- Leggett, S. K., et al. 2006, *MNRAS*, 373, 781
- Lehmer, B. D., et al. 2009, *ApJ*, 691, 687
- Lemaux, B. C., Lubin, L. M., Shapley, A., Kocevski, D., Gal, R. R., & Squires, G. K. 2010, *ApJ*, 716, 970
- Lidman, C., et al. 2008, *A&A*, 489, 981
- Magorrian, J., et al. 1998, *AJ*, 115, 2285
- Matsuda, Y., et al. 2004, *AJ*, 128, 569
- Matsuda, Y., et al. 2009, *MNRAS*, 400, L66
- Mei, S., et al. 2009, *ApJ*, 690, 42
- Moorwood, A. F. M., van der Werf, P. P., Cuby, J. G., & Oliva, E. 2000, *A&A*, 362, 9
- Nakata, F., Bower, R. G., Balogh, M. L., & Wilman, D. J. 2005, *MNRAS*, 357, 679
- Nordon, R., et al. 2010, *A&A*, 518, L24
- Onodera, M., Arimoto, N., Daddi, E., Renzini, A., Kong, X., Cimatti, A., Broadhurst, T., & Alexander, D. M. 2010, *ApJ*, 715, 385
- Oemler, A., Jr., Dressler, A., & Butcher, H. R. 1997, *ApJ*, 474, 561
- Osterbrock, D. E. 1989, *Astrophysics of gaseous nebulae and active galactic nuclei*, University Science Books
- Packham, C., et al. 2003, *MNRAS*, 345, 395
- Pannella, M., et al. 2009, *ApJ*, 701, 787
- Papovich, C., et al. 2007, *ApJ*, 668, 45
- Papovich, C., et al. 2010, *ApJ*, 716, 1503
- Patel, S. G., Holden, B. P., Kelson, D. D., Illingworth, G. D., & Franx, M. 2009, *ApJL*, 705, L67
- Pentericci, L., Roettgering, H. J. A., Miley, G. K., Carilli, C. L., & McCarthy, P. 1997, *A&A*, 326, 580
- Pentericci, L., et al. 2000, *A&A*, 361, L25
- Poggianti, B. M., Smail, I., Dressler, A., Couch, W. J., Barger, A. J., Butcher, H., Ellis, R. S., & Oemler, A., Jr. 1999, *ApJ*, 518, 576
- Poggianti, B. M., Bridges, T. J., Komiyama, Y., Yagi, M., Carter, D., Mobasher, B., Okamura, S., & Kashikawa, N. 2004, *ApJ*, 601, 197
- Poggianti, B. M., et al. 2006, *ApJ*, 642, 188
- Poggianti, B. M., et al. 2008, *ApJ*, 684, 888
- Rakos, K., Schombert, J., & Odell, A. 2008, *ApJ*, 677, 1019
- Rees, M. J. 1985, *MNRAS*, 213, 75
- Rieke, G. H., Alonso-Herrero, A., Weiner, B. J., Pérez-González, P. G., Blaylock, M., Donley, J. L., & Marcillac, D. 2009, *ApJ*, 692, 556
- Rodighiero, G. et al. 2010, *A&A*, in press (arXiv:1005.1089)
- Röttgering, H. J. A., van Ojik, R., Miley, G. K., Chambers, K. C., van Breugel, W. J. M., & de Koff, S. 1997, *A&A*, 326, 505
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Seymour, N., et al. 2007, *ApJS*, 171, 353
- Savaglio, S., et al. 2005, *ApJ*, 635, 260
- Shimasaku, K., et al. 2003, *ApJL*, 586, L111
- Sobral, D., Best, P. N., Smail, I., Geach, J. E., Cirasuolo, M., Garn, T., & Dalton, G. B. 2010, *MNRAS*, in press (arXiv:1007.2642)
- Stanford, S. A., Eisenhardt, P. R., & Dickinson, M. 1998, *ApJ*, 492, 461
- Steidel, C. C., Adelberger, K. L., Dickinson, M., Giavalisco, M., Pettini, M., & Kellogg, M. 1998, *ApJ*, 492, 428
- Steidel, C. C., Adelberger, K. L., Shapley, A. E., Pettini, M., Dickinson, M., & Giavalisco, M. 2000, *ApJ*, 532, 170
- Steidel, C. C., Adelberger, K. L., Shapley, A. E., Erb, D. K., Reddy, N. A., & Pettini, M. 2005, *ApJ*, 626, 44 & Miley, G. K. 2004, *A&A*, 428, 793
- Stockton, A., Canalizo, G., & Maihara, T. 2004, *ApJ*, 605, 37
- Stockton, A., & McGrath, E. 2007, *Cosmic Frontiers*, ASP Conference Series, Vol. 379, 122
- Stott, J. P., et al. 2010, *ApJ*, 718, 23
- Strazzullo, V., et al. 2010, *A&A*, 524, A17
- Sulentic, J. W., Rosado, M., Dultzin-Hacyan, D., Verdes-Montenegro, L., Trinchieri, G., Xu, C., & Pietsch, W. 2001, *AJ*, 122, 2993
- Sun, M., Donahue, M., & Voit, G. M. 2007, *ApJ*, 671, 190
- Suzuki, R., et al. 2008, *PASJ*, 60, 1347
- Tadaki, K., et al. 2010, *PASJ*, submitted.
- Tamura, Y., et al. 2009, *Nature*, 459, 61
- Tanaka, M., Kodama, T., Arimoto, N., Okamura, S., Umetsu, K., Shimasaku, K., Tanaka, I., & Yamada, T. 2005, *MNRAS*, 362, 268

- Tanaka, M., De Breuck, C., Venemans, B., & Kurk, J. 2010, A&A, in press (arXiv:1005.2253)
- Thomas, D., Maraston, C., Bender, R., & Mendes de Oliveira, C. 2005, ApJ, 621, 673
- Thomas, D., Maraston, C., Schawinski, K., Sarzi, M., & Silk, J. 2010, MNRAS, 404, 1775
- Tokunaga, A. T., & Vacca, W. D. 2005, PASP, 117, 421
- Tran, K.-V. H., Franx, M., Illingworth, G. D., van Dokkum, P., Kelson, D. D., Blakeslee, J. P., & Postman, M. 2007, ApJ, 661, 750
- Tran, K.-V. H., et al. 2010, ApJL, 719, L126
- Uchimoto, Y. K., et al. 2008, PASJ, 60, 683
- Urquhart, S. A., Willis, J. P., Hoekstra, H., & Pierre, M. 2010, MNRAS, 406, 368
- Valdés, J. R., Berta, S., Bressan, A., Franceschini, A., Rigopoulou, D., & Rodighiero, G. 2005, A&A, 434, 149
- van der Werf, P. P., Moorwood, A. F. M., & Bremer, M. N. 2000, A&A, 362, 509
- van Dokkum, P. G., Franx, M., Fabricant, D., Illingworth, G. D., & Kelson, D. D. 2000, ApJ, 541, 95
- van Dokkum, P. G., & Franx, M. 2001, ApJ, 553, 90
- Venemans, B. P., et al. 2002, ApJL, 569, L11
- Venemans, B. P., et al. 2005, A&A, 431, 793
- Venemans, B. P., et al. 2007, A&A, 461, 823
- Wallace, L., & Hinkle, K. 1997, ApJS, 111, 445
- Yagi, M., Komiyama, Y., Yoshida, M., Furusawa, H., Kashikawa, N., Koyama, Y., & Okamura, S. 2007, ApJ, 660, 1209
- Yamada, T., et al. 2001, PASJ, 53, 1119
- Yoshida, M., et al. 2002, ApJ, 567, 118
- Yoshikawa, T., et al. 2010, ApJ, in press (arXiv:1005.4727)
- Zirm, A. W., et al. 2008, ApJ, 680, 224







**Fig. 1. (Top)** The simulated shift of the transmission curve of the NB2289 filter. The horizontal axis is the wavelengths in  $\mu\text{m}$  and the vertical axis has arbitrary units. The curves with orange (dot-dot-dashed), red (thick solid), magenta (thick solid), and blue (thin solid) colors each indicate the transmission curves at  $x=256$  (close to the innermost FOV coverage), 1024 (detector center), 1536, and 2048 (outer edge of the FOV) in channel-2 raw-image coordinates (the bottom figure, and corresponds to the right-half of the MOIRCS FOV). At each  $x$  position, the shift of the transmission curve along with  $y$ -coordinates is also shown: the curve at the longest wavelength is at the detector center ( $y=1024$ ), and shifts to shorter wavelengths as the  $y$  coordinates goes to the edge of the detector ( $y=512/1536$ :middle and  $1/2048$ :bluest). The red arrow indicates the expected position of the  $\text{H}\alpha$  emission, and the vertical green lines (dots) indicate the range of the shift for objects at  $\pm 500 \text{ km s}^{-1}$  relative to the radio galaxy. The peak amplitude of the transmission curves at each  $x$  coordinate are artificially shifted for clarity. See discussion for §2.1.1 for details. **(Bottom)** The schematic figure of the MOIRCS field of view for explanation of the top figure. The full  $4' \times 7'$  FOV is divided into two channels. Note that the minimum  $x$  coordinate of the raw data area is around  $x = 270$  for channel 2 ( $x < 270$  is blocked by the mirror shadow). The curves on the channel-2 side show the iso-redshift contours (each number indicates the redshift) we can detect  $\text{H}\alpha$  at the central wavelength of the filter window. The NB2288 filter can cover the redshift range of  $\pm 0.017$ . The pointing center is around (270, 1024).



**Fig. 2.** The  $K_s$ -NB versus NB color-magnitude diagram for our cataloged objects (plus with the label “ALL”). The blue asterisks (labeled as “EM”) indicate our selected emission-line objects based on the criteria  $EW_o > 50 \text{ \AA}$ , the narrowband aperture magnitude of less than 22 mag, and  $\Sigma > 2$ , which is used for analysis in the paper (one spurious object in the sample is removed here). The brightest emitter is 4C 23.56 which is shown as a large asterisk. Emitters with red open boxes are the objects with MOIRCS spectra (except 4C 23.56). The tilted 4-dashed line indicates the 3-sigma limiting magnitude of  $K_s = 23.5$ . The insetted figure shows the transmission curve of the filters used in this study.




**Fig. 3.** MOIRCS MOS spectra of four H $\alpha$  emitters. The top spectra (HAE#491) is 4C 23.56. The barcode-like pattern at the bottom is the OH airglow spectra. The numbers at the bottom are wavelengths in angstrom.




**Fig. 4.** MOIRCS MOS spectra of four H $\alpha$  emitters. The top left spectra (HAE#491) is 4C 23.56. The OH airglow lines (red solid line) and the atmospheric transmission curve (blue dotted) are also plotted at the bottom panel of each spectra. Note that the spectra are not calibrated.







**Fig. 5. (a: left)** The  $B-J$  versus  $J-K_s$  two-color diagram for objects in the INGRID subfield. Five red boxes with the label “HAE(OKAY)” indicate the HAE candidates for which we can measure colors. Small plus symbols indicate other NB-selected objects in the area. The overplotted model tracks from blue to red indicate the “pure disk” model with continuous SFR (bluest), “50% disk + 50% bulge”, “20% disk + 80% bulge”, “pure bulge” (reddest, and mostly out of the figure) models, respectively. The cyan-colored region on each model track indicates the range  $2.0 < z < 3.8$ , and the farthest point in each cyan-colored track is around  $z = 2.5$  to  $3$ . See text for more details. **(b: right)** The same figure for galaxies in GOODS-North field (see Appendix for details). The data is from the MOIRCS Deep Survey (MODS) Catalog with  $K_s < 23$  mag (Kajisawa et al. 2009; Kajisawa et al. 2010). In the figure, red filled circles indicate objects with spectroscopic redshifts of  $2.3 < z < 2.6$ , similar redshifts to our H $\alpha$  emitters at  $z \sim 2.5$ . Blue triangles indicate objects at  $0.2 < z < 0.25$  (open) as well as  $0.75 < z < 0.8$  (filled), each of which corresponds to the redshift range for our foreground Paschen series (Pa $\alpha$  and Pa $\beta$ , respectively) interlopers, and the filled boxes (skyblue) indicate objects at  $3.3 < z < 3.8$  to which the background [O III] emitter contaminant for our data would belong. Inset histogram shows the fraction of objects with known spectroscopic redshift that satisfy the  $BJK$  criteria we used here as the object selection at  $z > 2$ . Note that there is a very limited number of spectroscopically confirmed galaxies at  $z > 2.7$ , as indicated by the large  $N^{1/2}$  errorbars.




**Fig. 6.** The distribution of the HAEs on the NB2288 image. The FOV covers  $7' \times 4'$  on the sky, with north and east to the top and left, respectively. The red open circles indicate the candidate HAEs by our selection. The three small open boxes indicate the Distant Red Galaxies from Kajisawa et al. (2006a) that show emission-line excess ( $K_s - \text{NB} > 1$ ) as seen in Fig.13 (also see Appendix). Two of them are also identified as HAEs. The remaining one is the only NB-excess object among the ERO sample by KC97. Two small open circles (green) indicate the candidate emission-line objects that satisfy the criteria of  $\Sigma > 2$  but are fainter than  $\text{NB}=22.0$ . Three red open circle with the letter "S" indicate the emitters with the MOIRCS spectra (§3.1.1). 4C 23.56 is shown as a large diamond. The area enclosed by the white box indicates the same area shown in Fig.7. The yellow rectangular region indicates the INGRID subfield (see §3.1).



**Fig. 7.** Zoom-up views of the “Eastern clump” region in both the NB2288 (right) and the  $K_s$  (left) images. The objects at the center of each yellow circle represent the HAEs. The  $1\sigma, 4\sigma, 7\sigma, 10\sigma, \& 13\sigma$  contours based on the rms noise of the  $K_s$  image before smoothing is applied to the Gaussian-smoothed  $K_s$  images, and is also overlaid on the NB2288 image. A clear extended or offset light distribution of HAEs in the NB2288 image is seen for (at least) the three objects on the right. Interestingly, the directions of the offsetted light distributions for these three objects are roughly the same (to the North-West). The size of the area is about  $26'' \times 26''$  ( $\sim 0.5 \times 0.5$  Mpc<sup>2</sup> comoving).




**Fig. 8.** The difference of the object position between the  $K_s$  and the NB2288 images. Objects that have similar magnitudes to our HAEs are used (dots). The 1-, 2-, and 3- $\sigma$  scatter of the distribution are shown as dashed circles in the figure from the inner to outer, respectively. The result for the HAEs is shown by the filled squares, with the 4 HAEs in eastern clump in Fig.7 overlaid by asterisks. The offset is shown in X and Y pixel coordinates, where the MOIRCS pixel scale is  $0.''117/\text{pixel}$  (corresponding physical scale is 0.96 kpc/pixel at  $z = 2.48$ ).




**Fig. 9.** The density contour of the faint *Spitzer* MIPS 24 $\mu$ m sources is overlaid on the MIPS 24 $\mu$ m image. The area with an exposure coverage less than 1/3 of the total is not shown here. The brightest source in the center is 4C 23.56. The red rectangular region indicates the MOIRCS field of view (North is up: see Fig. 6), and the red open circles indicate the HAEs. The distribution of the faint MIPS sources ( $90 < F_{24\mu\text{m}} [\mu\text{Jy}] < 275$ ) is smoothed with a Gaussian kernel with  $\sigma = 30''$ , and shown as contours. The contour interval is  $(2i + 1)$ -times ( $i = 0, 1, 3, 6, 11$ ) the square root of the average MIPS source density (from the “outer region”: see text) above average.







**Fig. 10.** The  $K_s$  versus stellar mass relation for objects at  $2.3 < z < 2.6$  from the MOIRCS Deep Survey (MODS) catalog used in Kajisawa et al. (2009). The catalog includes the compilation of the spectroscopic redshift data published, and are shown as crosses in the figure. The stellar mass is based on the SED-fitting by the 12-band photometry from UV to infrared, with the SED model from Bruzual & Charlot (2003). Details of the model fitting procedure can be found in Kajisawa et al. (2009).



**Fig. 11. (a: left)** The observed star-formation rate (SFR) of the HAE sample as a function of stellar mass. The plus marks indicate the  $\text{H}\alpha$ -based SFR *before* the extinction correction, while red filled box connected to each plus mark is its extinction-corrected SFR values (see text). The blue filled boxes are the SFR based on the *Spitzer* MIPS  $24 \mu\text{m}$  photometry. The dotted line (blue) indicates the mass-SFR relation at  $z \sim 2$  from Daddi et al. (2007). The two arrows indicate the objects with the upper-(lower-)limit for their mass (SFR before extinction correction) due to their faintness. The most massive object is 4C 23.56. Its MIPS-based SFR is not shown (due to an AGN contribution and the model limit). **(b: right)** The comparison of the  $\text{H}\alpha$ -based specific SFR (SSFR) of our HAE sample and of some field  $\text{H}\alpha$ -emitting objects as a function of stellar mass. The marks with dust extinction correction (red filled box) and raw values (plus) are connected by vertical lines. The field samples are from Förster-Schreiber et al. (2009) and Onodera et al. (2010). Note that Förster-Schreiber et al. (2009) shows the results using two types of the extinction correction; thus they are shown as connected vertical lines in the figure. Our SSFR estimate should be treated as the crude values due to the large uncertainty in both the extinction correction and the mass estimate. Note that 4C 23.56 is the point at  $M > 10^{12} M_{\odot}$ , and should be ignored because of strong AGN contamination in both the  $K$ -band and the  $\text{H}\alpha$  emission.



**Fig. 12.** Comparison of the  $K_s$  magnitude and  $L(H\alpha)$  for H $\alpha$  emitters from two field surveys from Förster-Schreiber et al. (2009) and Onodera et al. (2010) (asterisk and cross), PKS 1138-262 field from Kurk et al. (2004a) (box), and our study (plus). Only objects at  $z > 2$  are shown for field survey data. For the PKS 1138 field, we plot the data with the significance level over 3 and  $EW_o > 50\text{\AA}$  for better match to our selection criteria. Two arrows at brighter than  $K = 20$  represents the PKS 1138 and 4C 23.56 radio galaxies which are out of the figure due to their bright magnitudes.



**Fig. 13.** Color excess of objects studied by others in the 4C 23.56 field. Only objects identified in our catalog are shown here. The grey plus marks are all objects in the field of view, and the asterisks are the HAEs. A red filled box and 4 green open circles each indicate a Ly $\alpha$  emitter and the EROs from KC97 that are in our catalog. Blue filled circles indicate the DRG sample from Kajisawa et al. (2006b), and a magenta open triangle is the object studied by Stockton et al. (2004). Ignoring the radio galaxy, three DRGs/EROs show the emission-line excess ( $K_s - \text{NB} > 1$ ), and they are also plotted in Fig.6.



**Table 1.** Observational properties of candidate H $\alpha$  emitters.

HAE ID	RA(J2000)	DEC(J2000)	$K_s$ <sup>1</sup>	EW <sub>o</sub> <sup>2</sup>	F(H $\alpha$ ) <sup>3</sup>	F(24 $\mu$ m) <sup>4</sup>	redshift <sup>5</sup>
153	21:07:16.96	23:30:30.5	21.82 $\pm$ 0.07	327	3.64 $\pm$ 0.77	78.9	2.4879
354	21:07:21.03	23:31:13.9	21.23 $\pm$ 0.05	105	2.47 $\pm$ 0.76	134.9	2.4865
356	21:07:15.70	23:31:11.9	22.78 $\pm$ 0.12	282	1.35 $\pm$ 0.50	125.9	...
393 <sup>6</sup>	21:07:21.48	23:31:19.5	21.72 $\pm$ 0.06	51	0.81 $\pm$ 0.55	42.4	...
431	21:07:15.93	23:31:27.5	22.17 $\pm$ 0.08	221	1.96 $\pm$ 0.62	156.6	...
479 <sup>7</sup>	21:07:21.63	23:31:41.2	22.37 $\pm$ 0.08	956	4.25 $\pm$ 0.72	131.8	...
491 <sup>8</sup>	21:07:14.80	23:31:45.1	19.49 $\pm$ 0.02	122	13.95	4630	2.4852
500 <sup>9</sup>	21:07:21.84	23:31:41.8	> 23.5	> 780	> 0.8	...	...
511 <sup>10</sup>	21:07:22.58	23:31:43.4	22.98 $\pm$ 0.12	300	1.18 $\pm$ 0.46	156.5	...
526 <sup>11</sup>	21:07:21.78	23:31:49.6	20.76 $\pm$ 0.04	95	3.45 $\pm$ 0.94	119.2	2.4872
543	21:07:17.61	23:31:50.5	> 23.5	> 505	> 0.8	...	...

1. the SExtractor MAG\_AUTO output in AB mag.
2. rest-frame equivalent width based on the ( $K_s$ –NB2288) values by MAG\_AUTO output.
3. continuum subtracted narrowband flux in  $10^{-16}$  erg sec $^{-1}$  cm $^{-2}$ .
4. Spitzer MIPS 24  $\mu$ m flux in  $\mu$ Jy from the DAOPHOT photometry.
5. see §3.1.
6. a possible low redshift interloper based on the *BJK* criterion (§3.1).
8. 4C 23.56. Its  $K_s$  magnitude is uncertain due to stray light (see §2.1).
- 7,9,10,11. galaxies in the “Eastern Clump” (Fig.7; see §3.3).

**Table 2.** The estimated basic properties of the H $\alpha$  emitters.

HAE ID	log(M/M $_{\odot}$ )	$L_{H\alpha}$ <sup>1</sup>	SFR <sub>H<math>\alpha</math></sub> <sup>raw</sup>	SFR <sub>H<math>\alpha</math></sub> <sup>corr</sup> <sup>2</sup>	log( $L_{24 \mu m}$ /L $_{\odot}$ )	SFR(24 $\mu$ m) <sup>3</sup>
153	11.124 $\pm$ 0.235	14.4 $\pm$ 3.0	67.13	187.96	12.533	410
354	11.430 $\pm$ 0.225	9.75 $\pm$ 3.02	45.57	355.45	12.792	743
356	10.635 $\pm$ 0.260	5.34 $\pm$ 1.96	24.94	122.21	12.758	688
393	11.178 $\pm$ 0.230	3.19 $\pm$ 2.16	14.92	14.92	12.237	207
431	10.948 $\pm$ 0.243	7.74 $\pm$ 2.45	36.17	177.23	12.864	877
479	10.844 $\pm$ 0.240	16.75 $\pm$ 2.82	78.30	383.67	12.781	724
491 <sup>4</sup>	< 12.320 <sup>5</sup>	55.0 $\pm$ 6.8	257.15	1429.75 <sup>6</sup>	N/A <sup>7</sup>	N/A <sup>8</sup>
500	< 10.164	> 4.63	> 24.85	...	...	...
511	10.531 $\pm$ 0.262	4.65 $\pm$ 1.8	21.72	106.43	13.091	876
526	11.667 $\pm$ 0.220	13.6 $\pm$ 3.7	63.67	311.98	12.869	647
543	< 10.164	> 4.13	> 19.28	...	...	...

1. Extinction-uncorrected values in unit of  $10^{42}$  erg s $^{-1}$ .
2. in M $_{\odot}$  yr $^{-1}$ . See text for detail of the extinction correction.
3. *Spitzer* MIPS 24  $\mu$ m-based SFR in M $_{\odot}$  yr $^{-1}$ .
- 4,5. 4C 23.56. Its mass estimate is unreliable due to the uncertainty in its  $K_s$  magnitude (see §2.1). If we use the value by KC97, the estimated stellar mass is 11.67.
6. Affected by an AGN contribution. The SFR value is quite uncertain.
- 7,8. Affected by an AGN contribution. The  $L_{24 \mu m}$  conversion from the observed MIPS flux is impossible due to the limitation of the Chary and Elbaz (2001) model.